



## RESEARCH ARTICLE

### CALLUS INDUCTION AND SHOOT PROLIFERATION OF JASMINE (*JASMINUM SAMBAC* L. AITON.)

\*<sup>1</sup>Biswal, M., <sup>1</sup>Palai, S.K., <sup>1</sup>Mishra, P., <sup>2</sup>Chhuria, S., <sup>3</sup>Sahu, P.

<sup>1</sup>Department of Floriculture and Landscaping, College of Agriculture, Orissa University of Agriculture and Technology, Bhubaneswar-751003

<sup>2</sup>Department of Fruit Science and Technology, College of Agriculture, Orissa University of Agriculture and Technology, Bhubaneswar-751003

<sup>3</sup>Department of Vegetable Science, College of Agriculture, Orissa University of Agriculture and Technology, Bhubaneswar-751003

#### ARTICLE INFO

##### Article History:

Received 27<sup>th</sup> February, 2016  
Received in revised form  
19<sup>th</sup> March, 2016  
Accepted 22<sup>nd</sup> April, 2016  
Published online 10<sup>th</sup> May, 2016

##### Key words:

Bud break, Callus induction,  
MS medium,  
*Jasminum sambac*,  
Shoot Proliferation.

#### ABSTRACT

The present investigation was carried out on an important ornamental plant *Jasminum sambac* L. Aiton. belonging to family Oleaceae. The different vegetative parts i.e. nodal explants, shoot apices, stem and leaves were used to standardize the protocol through *in vitro*. The explants excised from field grown mature plant and thereafter planted on variously supplemented Murashige and Skoog's medium for multiple shoot proliferation and callus induction. Early bud break (25-26 days) and maximum percentage of bud break (83.7) was observed in *Jasminum sambac* when MS medium supplemented with BAP (2.0 mg/l), Kn (1.0 mg/l), Ads (50 mg/l). Maximum percentage of multiple shoots was observed in *Jasminum sambac* on MS medium fortified with 2 mg/l BAP, 1.0 mg/l Kn and 50 mg/l Ads. Callus induction from leaf and meristem explants were achieved using different concentrations of auxins and cytokinins. Maximum percentage of cultures showing callus proliferation from leaf explants was observed in the media containing MS basal salts supplemented with 3.0mg/l BAP and 2.0 mg/l IAA in leaf explants. The percentage of cultures showing callus was similar in the media containing MS basal salts supplemented with 5.0 mg/l BAP and 2.0 mg/l NAA in leaf explants. However, white and friable callus were obtained in the media containing MS basal salts supplemented with 2.0mg/l BAP, 2.0 mg/l Kn and 2.0 mg/l IAA from meristem explants.

Copyright © 2016, Biswal et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Biswal, M., Palai, S.K., <sup>1</sup>Mishra, P., Chhuria, S., Sahu, P., 2016. "Callus induction and shoot proliferation of jasmine (*Jasminum sambac* L. Aiton.)", *International Journal of Current Research*, 8, (05), 30164-30168.

## INTRODUCTION

Among commercial loose flowers, jasmine (*Jasminum sambac*) of the family oleaceae, native of India, is endowed with large spectrum of commercial potentialities in perfumery and essential oil sector. (Ambasta 1986; Chadha, 1978). The species is highly variable, possibly a result of Spontaneous mutation, Natural Hybridization, and Autopolyploidy. Only a few varieties are propagated by seed in the wild. Cultivated *Jasminum sambac* generally do not bear seeds, as a result, it is difficult to develop new varieties through conventional breeding. The only way to develop new cultivars is either through somatic mutations or gene manipulation.

Therefore, it is essential to develop an efficient protocol for *in vitro* culture of *Jasminum sambac*. Moreover, as it does not bear seeds, the plant is reproduced solely by cuttings and layering and these methods of propagation are dependent on the season. Layering involves more time and restricts the number of plants propagated from a bush. It can also be propagated by cuttings. However, long term cutting/layering causes varietal degeneration, resistance weakness and declining of flower production (Cai *et al.*, 2007).

*In-vitro* culture presents itself as an attractive tool for mass multiplication of important fragrant plant species and for the production of secondary metabolites. There are few reports on regeneration of jasmine employing tissue culture techniques (Jonard, 1989; Khoder *et al.* 1979). The present study will examine the efficiency of nodal buds in forming multiple shoots either with agar solidified medium or agar free liquid medium.

\*Corresponding author: Biswal, M

Department of Floriculture and Landscaping, College of Agriculture, Orissa University of Agriculture and Technology, Bhubaneswar-751003

## MATERIALS AND METHODS

### Plant material and explants preparation

An experiment was conducted at Agricultural Biotechnology Department, OUAT Bhubaneswar during the year 2013-14. stem apices and leaves were used as source of explants. Young shoots were selected from garden. the shoot materials either apical or axillary bud were first washed for 15 minutes in running tap water, cut into 1.5-2 c.m. then treated with 0.1% Bavistin solution for 15 minutes, followed by through washing (5-6 times) with distilled water. surface sterilization was carried out with 0.1%HgCl<sub>2</sub> (w/v) for 10 minutes and followed by washing 3 times with sterile distilled water.

### Culture media and culture conditions

For shoot initiation medium each test tube(150mmX25mm) with 10-15 ml of Murashige and Skoog (MS medium supplemented with 30% sucrose and various combination of BAP(1.0-6.0 mg/l), NAA (0.5-2 mg/l) and Ads(0-50mg/l) The medium containing 3% (w/v) sucrose and 0.7-0.8% (w/v) agar. For callus culture media was supplemented with various concentration of BA (1-5mg/l) or IAA (1-2mg/l) or NAA (1-2mg/l). The P<sup>H</sup> was adjusted to 5.7 with 1N KOH or 1N HCL before autoclaving at 1.05kg/cm<sup>2</sup>, 121° C for 20 minutes.

The explants (shoot tip) were cut into pieces (3-4cm), each containing one or two axillary buds and inoculated in the fortified medium. The leaf pieces placed to previously prepared slants. The cultures were incubated in growth chamber (25±2°C) under cool white flurorescent lamps providing 3000 lux light intensity. The photoperiod was adjusted to 16/8 h light and dark cycle. However for callus culture from leaf, The cultures were maintained at dark condition till sub culturing. Each treatment was represented by 20 explants and the experiment was repeated three times.

## RESULTS AND DISCUSSION

### Effect of Growth Regulators on Bud break

Bud break was achieved on MS medium supplemented with various concentrations of BAP, Kn, IAA and Ads. BAP and Kn favoured bud break among the different hormones tested. Adenine sulphate helped in inducing bud break at low concentration in combination with BAP and Kn. The auxins were less effective in inducing bud break than cytokinins. Early bud break (25-26 days) was initiated in *Jasminum sambac* when MS medium was supplemented with BAP (2 mg/l), Kn (1 mg/l) and Ads (50 mg/l) (Table 1). The maximum percentage (83.7) of bud break in *Jasminum sambac* was on MS medium

**Table 1. Effect of auxin and cytokinins on bud break of *Jasminum samabc* (L.) Aiton, grown in MS medium, after 8 weeks of culture**

MS + growth regulators (mg/l)				Days to bud break	Percentage of bud break (Mean ± S.E.)*
Kn	BAP	IAA	Ads		
0	0	0	0	0	0
0	1.0	0.1	50	34-36	4.8 ±0.4
0	2.0	0.1	50	35-37	13.3±0.7
0	3.0	0.1	50	36-38	21.6±0.5
1.0	0	0.1	50	35-38	10.0±2.4
2.0	0	0.1	50	33-36	12.0±1.3
3.0	0	0.1	50	44-47	15.6±1.2
1.0	1.0	0	50	26-28	76.6±1.4
1.0	2.0	0	50	25-26	83.7±1.8
1.0	3.0	0	50	25-27	77.6±1.3
2.0	1.0	0	50	31-33	75.0±1.1
2.0	2.0	0	50	37-39	73.3±1.4
2.0	3.0	0	50	32-34	76.6±1.7
3.0	1.0	0	50	38-40	61.6±1.4
3.0	2.0	0	50	40-42	62.4±1.5
3.0	3.0	0	50	36-38	54.8±1.6

\* 20 cultures per treatment; repeated thrice.

**Table 2. Effect of Auxin and cytokinins on shoot multiplication of *Jasminum sambac* (L.) Aiton, after 8 weeks of culture**

MS + growth regulators (mg/l)				Percentage of explants developed multiple shoot (Mean ± S.E.)*	Number of shoots/culture (Mean ± S.E.)*
Kn	BAP	IAA	Ads		
0	0	0	0	0	0
0	1.0	0.25	50	14.2 ±0.6	0.8 ±0.06
0	2.0	0.25	50	23.3±0.8	1.3±0.07
0	3.0	0.25	50	31.6±1.5	2.1±0.05
1.0	1.0	0	50	76.2±1.2	6.6±0.4
1.0	2.0	0	50	88.4±1.7	8.7±0.8
1.0	3.0	0	50	81.6±1.1	5.6±0.3
2.0	1.0	0	50	79.0±1.0	5.0±0.2
2.0	2.0	0	50	78.4±1.1	3.3±0.04
2.0	3.0	0	50	80.2±1.2	4.6±0.07
3.0	1.0	0	50	66.4±0.9	2.6±0.08
3.0	2.0	0	50	67.6±0.9	2.4±0.04
3.0	3.0	0	50	64.8±0.8	1.8±0.09

having BAP (2 mg/l), Kn (1 mg/l), Ads (50 mg/l) and 3% sucrose (Table 1). Depending on the type and concentration of growth regulators used, the number of days required to bud break varied between 25 to 47 days in *Jasminum sambac*. Among all growth regulators tested, the media containing BAP (2.0 mg/l), Kn (1.0 mg/l) and Ads (50.0 mg/l) induced bud sprouting within 25-26 days of culture of *Jasminum sambac*. BAP was reported to be in general, the most effective cytokinin for meristem, shoot-tip and axillary bud culture of various species (Cai *et al.*, 2007; Bhattacharya and Bhattacharya, 1997).

### Effect of Plant Growth Regulators on Shoot Multiplication

The experiment was designed to study the effect of different concentration and combinations of cytokinins (BA and Kinetin) and Auxins(IAA) on shoot initiation and elongation. The shoots showed differential rate of elongation. Media devoid of growth regulator did not respond. Maximum percentage of multiple shoot (88.4) was observed in MS medium supplemented with BAP (2.0 mg/l), Kn (1.0 mg/l) and Ads (50 mg/l) within 8 weeks of culture.

**Table 3. Effect of Liquid media on shoot multiplication of *Jasminum sambac* (L.) Aiton cultured on MS basal medium supplemented with 1.0 mg/l Kn , 2.0 mg/l BAP, 50 mg/l Ads after 8 weeks of culture**

MS + growth regulators (mg/l)				Percentage of explants developed multiple shoot (Mean $\pm$ S.E.)*		Number of shoots/culture (Mean $\pm$ S.E.)*	
Kn	BAP	IAA	Ads	Agar Media (0.8%)	Liquid media	Agar Media (0.8%)	Liquid media
0	0	0	0	0	0	0	0
0	1.0	0.25	50	12.5 $\pm$ 0.7	14.8 $\pm$ 0.9	0.6 $\pm$ 0.07	1.1 $\pm$ 0.3
0	2.0	0.25	50	21.5 $\pm$ 1.2	25.6 $\pm$ 1.5	1.4 $\pm$ 0.08	1.8 $\pm$ 0.6
0	3.0	0.25	50	32.9 $\pm$ 1.9	36.7 $\pm$ 1.8	2.3 $\pm$ 0.07	2.9 $\pm$ 0.8
1.0	1.0	0	50	73.6 $\pm$ 1.8	75.2 $\pm$ 1.4	5.6 $\pm$ 0.6	6.2 $\pm$ 0.9
1.0	2.0	0	50	86.9 $\pm$ 1.9	90.5 $\pm$ 1.2	8.6 $\pm$ 0.7	8.9 $\pm$ 0.9
1.0	3.0	0	50	79.1 $\pm$ 1.4	82.2 $\pm$ 1.9	4.9 $\pm$ 0.5	5.1 $\pm$ 0.7
2.0	1.0	0	50	76.2 $\pm$ 1.1	78.3 $\pm$ 1.6	4.6 $\pm$ 0.7	4.8 $\pm$ 0.3
2.0	2.0	0	50	75.6 $\pm$ 1.3	77.1 $\pm$ 1.9	3.0 $\pm$ 0.6	3.3 $\pm$ 0.4
2.0	3.0	0	50	78.9 $\pm$ 1.5	79.6 $\pm$ 1.2	4.1 $\pm$ 0.5	4.8 $\pm$ 0.4
3.0	1.0	0	50	68.6 $\pm$ 1.8	71.6 $\pm$ 1.5	2.2 $\pm$ 0.06	2.6 $\pm$ 0.09
3.0	2.0	0	50	65.2 $\pm$ 1.7	67.1 $\pm$ 1.2	2.0 $\pm$ 0.03	2.4 $\pm$ 0.07
3.0	3.0	0	50	61.7 $\pm$ 1.5	62.9 $\pm$ 1.3	1.5 $\pm$ 0.07	1.9 $\pm$ 0.05

\* 20 cultures per treatment; repeated thrice.



**Figures 1. In vitro propagation of *Jasminum sambac*. A. Enlarged axillary buds of jasminum sambac at the nodes after 2 weeks of culture. B. Development of single axillary shoot from nodal explants on MS medium supplemented with MS medium having BAP (2 mg/l), Kn (1 mg/l), Ads (50 mg/l) and 3% sucrose after 8 weeks of culture. C. Development of multiple shoots from nodal explants on liquid medium supplemented with BAP (2 mg/l), Kn (1 mg/l), Ads (50 mg/l) and 3% sucrose after 8 weeks of culture, D. induction of callus from leaf explants E. induction of callus from meristem explants**

Induction of multiple shoot was better on MS medium supplemented with BAP + Kn than BAP alone. IAA was less effective in inducing multiple shoot in combination with BAP and Ads. The average number of shoots per culture was the maximum (8.7) in MS medium supplemented with BAP (2.0 mg/l), Kn (1.0 mg/l) and Ads (50 mg/l) (Table 2).

BAP (2.0 mg/l), Kn (1.0 mg/l) and Ads (50 mg/l) (Table 3). In the present investigation, the rate of shoot multiplication in liquid medium (medium devoid of gelling agent) was better than the agar gelled medium.

**Table 4. Effect of growth regulator on callus induction in Meristem Explants and Leaf Explants of *Jasminum sambac* (L.) Aiton cultured after 8 weeks of culture**

MS+growth regulators (mg/l)			Percentage of Cultures showing callus	MS+ growth regulators (mg/l)			Percentage of cultures showing callus
			Meristem callus				Leaf callus
BAP	Kn	IAA		BAP	IAA	NAA	
0	0	0	0	0	0	0	0
1.0	1.0	1.0	32.5±1.4	1.0	1.0	0	12.3±0.4
2.0	1.0	1.0	38.4±1.2	2.0	1.0	0	18.4±0.6
3.0	1.0	1.0	27.3±0.8	3.0	1.0	0	37.6±0.7
1.0	2.0	1.0	27.7±0.3	4.0	1.0	0	27.7±0.3
2.0	2.0	1.0	42.6±1.2	5.0	1.0	0	22.6±0.6
3.0	2.0	1.0	29.5±0.7	6.0	1.0	0	23.5±0.8
1.0	3.0	1.0	36.6±0.9	1.0	2.0	0	33.5±0.4
2.0	3.0	1.0	52.7±0.7	2.0	2.0	0	56.7±0.5
3.0	3.0	1.0	36.6±0.8	3.0	2.0	0	86.7±1.8
1.0	1.0	2.0	22.7±0.4	4.0	2.0	0	52.3±0.4
2.0	1.0	2.0	39.4±0.8	5.0	2.0	0	38.4±0.6
3.0	1.0	2.0	42.2±0.9	6.0	2.0	0	34.2±0.7
1.0	2.0	2.0	67.8±1.4	1.0	3.0	0	27.8±0.4
2.0	2.0	2.0	89.6±1.5	2.0	3.0	0	39.6±0.5
3.0	2.0	2.0	55.1±0.8	3.0	3.0	0	52.1±0.9
1.0	3.0	2.0	35.7±0.7	4.0	3.0	0	46.7±0.8
2.0	3.0	2.0	45.3±0.8	5.0	3.0	0	47.3±0.8

The results indicate that rate of shoot multiplication of *Jasminum sambac* declined as the concentration of BAP increased from 2.0 to 3.0 mg/l. this may be due to the ionic concentration and balance between different nutrient ions in the culture medium which is crucial for optimization of shoot multiplication of *Jasminum sambac*. Similar observations were made on *Jasminum sambac* by Cai *et al.*, (2007) and Sun *et al.*, (2009). The present study indicated that inclusion of IAA in the medium, either with BAP or Kn, did not favour multiplication and growth of the micro shoots. The cytokinin requirement for shoot multiplication was essential and two cytokinins (BAP and Kn) favoured shoot multiplication. However, Cai *et al.* (2007) reported that MS medium in combination with BA (2.0 mg/l) + NAA (1.0 mg/l) and MS medium in combination with BA (1.5 mg/l) + NAA (0.3 mg/l) + GA<sub>3</sub> (0.5 mg/l) favoured axillary bud initiation and axillary bud proliferation respectively.

#### Effect of Liquid Media on shoot multiplication

The effect of Liquid media on shoot multiplication was studied in comparison with agar media (0.8%). Twenty milliliter of liquid media was taken in each test tube and absorbent cotton/filter paper bridges were inserted in the tubes prior to autoclaving. The absorbent cotton/filter paper bridges helps the explants to remain in proper position. Induction of multiple shoot was better on MS liquid medium supplemented with BAP + Kn than BAP alone. In *Jasminum sambac* maximum percentage of multiple shoot (90.5) was observed in MS liquid medium supplemented with BAP (2.0 mg/l), Kn (1.0 mg/l) and Ads (50 mg/l). The average number of shoots per culture was the maximum (8.9) in MS liquid medium supplemented with

This may be due to easier and efficient translocation of nutrients in the liquid medium than agar gelled medium. Similar results were reported in other species of jasmine (Dainty *et al.*, 1985).

#### Effects of Media supplement on Callus Induction and Regeneration

Proliferation in leaf explants and callus induction were achieved by using MS medium supplemented with different auxins (IAA or NAA) and cytokinin (BAP) alone or in combinations. Calli were initiated from leaf explants on the MS basal medium supplemented with BAP + NAA or BAP + IAA within 13 days of inoculation. The initial calli were small, globular and pale yellow in colour which developed on the surface of the explant and subsequently spread over the entire explant. The maximum percentage (86.7) of cultures showing callus in leaf explants was observed in MS medium supplemented with BAP (3.0mg/l) and IAA (2.0mg/l) which was at par with the MS medium supplemented with BAP (5.0mg/l) and NAA (2.0mg/l).

The differential response could be due to the varying concentrations of growth regulators used in the media. Similar observations have been reported for *Jasminum grandiflorum* (Gomathi *et al.*, 2007). *Jasminum malabathricum* Wight. (Gadkar *et al.*, 2011) and in *Jasminum sambac* (Nurmalita *et al.*, 2012). However, Induction of callus and its proliferation by using meristem explants of *Jasminum sambac* was achieved on the MS basal medium supplemented with BAP, Kn and IAA within 21 days of inoculation which were small, globular and white in colour. Maximum callus growth was obtained in the

media containing MS basal salts supplemented with 2.0 mg/l each of BAP, Kn and IAA. The calli were subsequently transferred to different MS medium supplemented with various concentration of cytokinins and auxins for plant regeneration. However, the shoot bud like structures have also been achieved in medium having MS medium supplemented with 3.0 mg/l kinetin or BAP (2-3mg/l) in combination with IAA or NAA (0.5-3.0 mg/l). Beasley and Pijut (2013) reported that the plant regeneration from leaf and stem explants of *Fraxinus nigra* by using the medium of 2.5 mg/l BAP and 0.5 mg/l NAA.

### Conclusion

High frequency plant regeneration of *Jasminum sambac* through *in vitro* techniques was attempted by manipulation of nutrient media and culture conditions. Apical and axillary meristems were used as explant sources for shoot multiplication. The frequency of shoot multiplication was the highest in MS medium supplemented with BAP, Kn and Ads. Bud break was achieved on MS medium supplemented with BAP, Kn, IAA and Ads in *Jasminum sambac*. Early bud break (25-26 days) was observed in *Jasminum sambac* when MS medium was supplemented with BAP (2.0 mg/l), Kn (1.0 mg/l), Ads (50 mg/l). Maximum percentage of bud break and multiple shoot formation (83.7, 90.5) in *Jasminum sambac* was observed on MS medium having BAP (2.0 mg/l), Kn (1.0 mg/l), Ads (50 mg/l) and 3% sucrose within 4 week of subculture.

In comparisons between solid agar gel medium with liquid medium, it was observed that the shoot multiplication in liquid medium was found to be better than agar gelled medium. Callus induction in leaf and meristem explants were achieved using different concentrations of auxins and cytokinins. Maximum percentage of cultures showing callus in *Jasminum sambac* was observed in the media containing MS basal salts supplemented with 3.0 mg/l BAP and 2.0 mg/l IAA in leaf explants, Which was at par in the media containing MS basal salts supplemented with 5.0 mg/l BAP and 2.0 mg/l NAA in leaf explants. However, white and friable callus were obtained in the media containing MS basal salts supplemented with 2.0 mg/l BAP, 2.0 mg/l Kn and 2.0 mg/l IAA in meristem explants. Further work is necessary to achieve the plant regeneration from proliferated calli to sustain the commercial propagation. Several combinations of auxins and cytokinin were tried to regenerate shoots from the callus and induce rooting in micro-shoots of *Jasminum sambac* but, it could not be achieved during the period of study. However, the protocol developed in the present study can be taken up as a base for future experiment on plant regeneration of *Jasminum sambac*.

### REFERENCES

- Ambasta, S.P. 1986. In: Ambasta SP (ed) the useful plants of india (pp301). Publication and information Directorate, CSIR, New Delhi
- Beasley, R. R. and Pijut, P. M. 2013. Regeneration of plants from *Fraxinus nigra* Marsh. Hypocotyls. *Hort Science*, 48 (7): 887-890
- Bhattacharya, S. and Bhattacharya, S. 1997. Rapid multiplication of *Jasminum officinale* L. by *in vitro* culture of nodal explants. *Plant Cell, Tissue and Organ Culture*, 51:57-60.
- Bhattacharya, S and Bhattacharyya, S. 2010. *inVitro* Propagation of *Jasminum officinale* L. A Woody Ornamental Vine Yielding Aromatic Oil from Flowers. *Protocols for In Vitro Propagation of Ornamental Plants; Methods in Molecular Biology*, Volume 589: 117-126
- Cai, H., Chen, X., Xiong, Z., Xie, L. and Zhao, L. 2007. Techniques of *in vitro* micro-propagation and sugar-free rooting of Jasmine (*Jasminum sambac*). *Jiangsu Journal of Agricultural Sciences*, 23(5):464-468.
- Chadha YR (1976) in : Chadha YR (ed) the wealth of india, raw materials(pp279). publication and information Directorate, CSIR, New Delhi
- Dainty, A.L., Goulding, K.H., Robinson, P.K., Simpkins, I. and Trevan, M.D. 1985. Effect of immobilization on plant cell physiology real or imaginary trends. *Biotechnology* 3(3):59-60.
- EKA Nurmalita-Sari, Nurmalita sari yulia, Lukas Suhendra Budipramana and Evie Ratnasari, 2012. Callus Induction and Growth of Stem Melati (*Jasminum sambac*) on MS medium with the addition of Gibberellins. *Lantera Bio* Vol.1, no.1
- Rout, G. A. Mahato and S. Senapati, 2008. *In vitro* clonal propagation of *Nyctanthes arbor-tristis*. *Biologia Plantarum.*, Vol. 52 (3): 521-524
- Gadkar *et al.* 2011. tissue culture studies on *Jasminum malbaricum* wight. for Micropropagation. Department of pharmaceutical biotechnology KLEU's college of pharmacy Belgaum.
- Gomathi, K. S., Sambandamurthy, S., Sadasivam, S., Ramasamy, N. M. and Rajmohan, K. 2007. *In vitro* culture and biochemical analysis of *Jasminum* sp. In: *Recent trends in Horticultural Biotechnology*, Vol. I and II.
- Khoder, M.; Villemur, P. and Jonard, R. 1979. *In vitro* micropropagation and stem cuttings of *Jasminum officinale*. *Comptes Rendus Hebdomadaires des Seances de l'Academie des Sciences*, 228 (3): 323-326
- Sun, Y.; Tang, F.; Fang, W. and Tan, G. 2009. Rapid propagation *in vitro* of jasmine. *Acta Agriculturae Zhejiangensis*, 21(4):390-394.

\*\*\*\*\*