



FREE RADICAL SCAVENGING AND ANTI-LIPIDPEROXIDATIVE POTENTIAL OF *Calocybe indica*

¹S. Selvi #, ¹J. Dhanalakshmi, ²P. Umadevi and ³D. Devipriya

¹Department of Biochemistry, Bharathidasan College of Arts and Science, Erode

²Department of Biotechnology, Sri Narayanaguru Arts and Science College, Coimbatore

³Dr. N.G.P Arts and Science College Coimbatore

ARTICLE INFO

Article History:

Received 24th March, 2012

Received in revised form

27th April, 2012

Accepted 25th May, 2012

Published online 30th June, 2012

Key words:

Free radical scavenging,

Calocybe indica,

Antioxidant.

ABSTRACT

The ethanolic extract of *Calocybe indica* was studied for its free radical scavenging property on different *in vitro* models like 1,1-diphenyl-picrylhydrazyl (DPPH) Assay, Ferric Reducing Antioxidant Power (FRAP) assay and *in vitro* antilipidperoxidative assay using goat liver homogenate and RBC Ghost model. The *in vitro* Lipid peroxidation (LPO) was inhibited to a good extent by the *Calocybe indica* ethanolic extract and the extent of inhibition being higher in the RBC membrane model than the liver homogenate model. The mushroom extract showed good dose-dependent free radical scavenging property in both the models.

Copy Right, IJCR, 2012, Academic Journals. All rights reserved.

INTRODUCTION

The study of lipid peroxidation (LPO) is attracting much attention in recent years due to its role in disease processes. Membrane lipids are particularly susceptible to LP due to the presence of polyunsaturated fatty acids. Since membranes form the basis of many cellular organelles like mitochondria, plasma membrane, endoplasmic reticulum, lysosomes, peroxisomes, etc. the damage caused by LP is highly detrimental to functioning of the cell and its survival. It has been implicated in the pathogenesis of a number of diseases and clinical conditions. These include atherosclerosis, cancer, adult respiratory distress syndrome, Alzheimer's disease, Parkinson's disease, ischaemia-reperfusion injury of various organs, chemical and radiation-induced injury, diabetes, etc. Experimental and clinical evidence suggests that aldehyde products of LP can also act as bioactive molecules in physiological and pathological conditions¹.

The most deleterious effect caused by ROS (Reactive oxygen species) is the peroxidation of membrane lipids. Exogenous chemicals and radiation produce peroxidation of lipids leading to structural and functional damage to cellular membranes². Polyunsaturated fatty acids present in cellular membranes are especially prone to damage by ROS and the resulting LP can have serious consequences. LP plays a major role in mediating oxidative-damage in biological systems. There are also several toxic by-products of peroxidation which can damage other biomolecules away from the site of generation^{3,4}. Antioxidants may offer resistance against oxidative stress by scavenging the free radicals, inhibiting lipid peroxidation.

Among the fungi, mushrooms have been used for untold centuries as food and medicine. Edible and medicinal mushrooms not only convert the huge lignocellulosic biomass waste into human food; but most remarkably, can produce notable myco-pharmaceuticals, myco-nutriceuticals and myco-cosmeceuticals. Mushroom accumulates a variety of secondary metabolites, including phenolic compounds, polyketides, terpenes and steroids. Also, a mushroom phenolic compound has been found to be an excellent anti-oxidant and synergist that is not mutagenic⁵.

Epidemiological studies show that a high intake of anti-oxidant-rich foods is inversely related to cancer risk. Animal and Cell Cultures confirm the anticancer effects of antioxidants. Selenium and vitamin E reduced the risk of some forms of cancer, including prostate and colon cancer and carotenoids have been shown to help reduce breast cancer risk⁶. Cultivation of milky mushrooms has become popular in Tamil Nadu, Kerala, Karnataka and Andhrapradesh. This is a tropical mushroom. It is attractive milky white mushrooms with excellent shelf life. Since the role of free radicals has been implicated in large number of diseases, the antioxidant activity of mushrooms is of significant importance in exploiting their therapeutic potential⁷. At the molecular level, researchers have found that antigenotoxic factors in the mushrooms include polysaccharides, such as beta- and alpha-glucan⁸ which acts against cancer. Mushrooms are regarded as healthy foods and therefore they have become an interesting subject for research⁹.

*Corresponding author: ramselvi@rediffmail.com;

MATERIALS AND METHODS

Collection and Preparation of Sample: Mushroom samples were collected from Blue hill Mushroom Producers and the sample is been preserved in the Department of Biochemistry Dr N.G.P Arts and Science College, Coimbatore. The mushrooms were shade dried and made into a coarse powder. The coarse powder was extracted using ethanol for 72 hours in a Soxhlet's apparatus. The ethanol was evaporated and the extract was concentrated and was used for the assay.

Preparation of goat liver homogenate: Fresh goat liver was obtained from local slaughter house, washed free of blood and removed fat deposits, if any, a 5% homogenate was prepared in ice cold TBS (Tris Buffered Saline) and used for assay. The assay procedure given by Okhawa *et al.*, (1979)¹⁰ have been followed.

Table 1. Extent of Inhibition of *in vitro* Lipid Peroxidation in RBC Ghost and Goat Liver Homogenate by Ethanolic Extract of *Calocybe indica*

S.No	Concentration of Extract in(µg)	Percentage Inhibition	
		RBC Ghost	Liver Homogenate
1.	20	33.60 ± 0.11	10.20 ± 0.21
2.	40	42.24 ± 0.01	14.42 ± 0.11
3.	60	54.80 ± 0.21	28.24 ± 0.03
4.	80	63.41 ± 0.10	43.32 ± 0.01
5.	100	79.28 ± 0.04	56.25 ± 0.09

Values represented as mean ± SD: n= 3.

Table 2. *In vitro* Antioxidant Capacity of Ethanolic Extract Of *Calocybe indica* (DPPH assay)

S.No	Concentration of Extract in(µg)	Percentage Inhibition
1.	20	19.03 ± 0.12
2.	40	36.04 ± 0.01
3.	60	52.40 ± 0.11
4.	80	68.02 ± 0.12
5.	100	85.41 ± 0.03

α- tocopherol IC 50 = 18.2 µg

Values represented as mean ± SD: n= 3.

Table 3. *In vitro* Antioxidant Capacity of Ethanolic Extract of *Calocybe indica* (FRAP Assay)

S.No	Concentration of Extract in(µg)	Percentage inhibition
1.	20	0.147 ± 0.10
2.	40	0.220 ± 0.04
3.	60	0.288 ± 0.01
4.	80	0.368 ± 0.03
5.	100	0.428 ± 0.01

Values represented as mean ± SD: n= 3.

Preparation of Erythrocyte Ghosts: About 50ml of fresh venous whole blood of goat was collected into a cleaned sterile bottle and defibrinated immediately using acid-washed stones. The defibrinated blood was then transferred into sterile centrifuge tubes and spun at 3000 rpm for 10min to pellet out the cells and the supernatant was discarded. The pellet of RBCs was washed in isotonic TBS, thrice successively. The washed pellet was then treated with hypotonic TBS and incubated at 37°C for 1 hour for lysis to occur. The lysate was centrifuged at 5000 rpm for 15-20 min at 4°C. The pale pellet containing the erythrocyte ghost membranes' was then suspended in 1.5ml of TBS. The assay procedure given by Dodge *et al.*, (1963)¹¹ has been followed to study the anti-

lipidperoxidative effect of the mushroom. 1-Diphenyl 1-2 Picrylhydrazyl (DPPH).

Radical Scavenging Activity: The hydrogen atom or electron donation abilities of the corresponding extracts were measured from the bleaching of the purple-coloured methanol solution of 1, 1- Diphenyl-2-picrylhydrazyl (DPPH). This spectrophotometric assay uses the stable radical DPPH as a reagent (Brand-Williams 1999)¹². 2 to 1 ml of various concentrations of the ethanolic mushroom extract was added to 4 ml of 0.004% methanol solution of DPPH. After a 30 min incubation period at room temperature, the absorbance was read against a blank at 517 nm. Inhibition of free radical by DPPH in percent (I %) was calculated as follows:

$$I (\%) = (A \text{ blank} - A \text{ sample} / A \text{ blank}) \times 100$$

Where A blank is the absorbance of the control reaction (containing all reagents except the test compound), and A sample is the absorbance of the test compound. Extract concentration providing 50% inhibition (IC50) was calculated from the plot of inhibition (%) against extract concentration. Tests were carried out in triplicate.

Ferric Reducing Ability Of Plant (FRAP) as Measuring Of Antioxidant Power

FRAP assay was carried out by the method of (Pulido 2000)¹³ with minor modification. The method is based on the reduction of a ferric 2, 4, 6-tripyridyl-s-triazine complex (Fe³⁺-TPTZ) to the ferrous form (Fe²⁺-TPTZ). The ethonal extract of mushroom in varying concentration (20-100µg) were added to 10mM ferric-TPTZ reagent and the increase in absorbance at 593 nm was measured at 8 min.

RESULTS

Determination of *in vitro* LPO: The results obtained are shown in Table 1. The extracts tested were effective in reducing the production of TBARS in a dose –dependent manner. The *in vitro* LPO was inhibited to a good extent by the *Calocybe indica* extract and the extent of inhibition being higher in the RBC membrane model than the liver homogenate model.

Determination of *in vitro* Antioxidant assay (DPPH and FRAP): Since the role of free radicals has been implicated in a large number of diseases, the antioxidant activity of mushroom is of significant importance in exploiting their therapeutic potential. The ethanolic extract was subjected to screening for their possible antioxidant activity. Two complementary test system, namely DPPH Free Radical Scavenging and FRAP assay were used for the analysis. Results are indicated in the Table 2 and 3.

DISCUSSION

The scientific community, in searching for new therapeutic alternatives, has studied many kinds of mushrooms and found variable therapeutic activity such as anticarcinogenic, anti-inflammatory, immunosuppressor and antibiotic¹⁴. The Lipid peroxidation (LPO) has been broadly defined as the oxidative deterioration of polysaturated lipids. Peroxyl and hydroxyl

radicals are important agents that mediate lipid peroxidation, thereby damaging cell membranes¹⁵. A number of toxic compounds are generated during this process of LPO. TBARS are produced as by-products of LPO that occurs in the hydrophobic core of biomembranes¹⁶. A substance may act as an antioxidant due to its ability to reduce ROS by donating hydrogen atom¹⁷. Two different model systems namely goat liver homogenate and RBC ghost (plasma membrane) were used to compare the membrane models which differ in their lipid composition. Mammalian cells have evolved migrate interrelated antioxidant defense mechanisms, which minimize the injurious events that result from toxic chemicals and normal oxidative products of cellular metabolism¹⁸. The effect of mushroom extracts on LP show significant inhibition of TBARS formation. The present finding strongly suggests that the use of the mushroom extracts to prevent LP leading to membrane damage consequent to radiation and to certain chemicals which generate potent ROS.

DPPH, a stable free radical with a characteristic adsorption at 517 nm, was used to study the radical scavenging effects of the extracts. As antioxidants donate protons to these radicals, the absorption decreases. The decrease in absorption is taken as a measure of the extent of the radical scavenging. All the concentration which was studied showed free radical scavenging activity. The 50% of inhibition value for *Calocybe indica* (IC₅₀ = 43.05 µg ethanolic extract was necessary to obtain 50 % of DPPH degradation). Ethanolic extract seems to be fairly significant when compared to commonly used synthetic antioxidant alpha-tocopherol (α-tocopherol IC₅₀ = 18.2 µg). The inhibition value increases with increase in concentration. Current research is now directed towards finding naturally occurring antioxidants which could reduce or minimize the deleterious effect caused by ROS. The result of the investigation reveal that ethanolic extract of *Calocybe indica* have potent lipid peroxidation inhibition and showed the highest antioxidant capacity when determined by DPPH and FRAP assays.

REFERENCES

1. Devasagayam T PA, Bolor KK and Ramasarma T (2003) Methods for estimating lipid peroxidation: An analysis of merits and demerits (mini review). Indian J. Biochem. Biophys., 40.
2. Rayleigh J A, (1987) In Prostaglandin and Lipid Metabolism in Radiation Injury (eds Walden Jr. T.L and Hughes, H.N) Plenum Press. Newyork vol 3,8.
3. Esterbauer H, (1996) Estimation of peroxidative-damage: A critical review. Pathol. Bio Paris, 44, 25–28.
4. Box HC, and Maccubbin AE (1997) Lipid peroxidation and DNA damage. Nutrition, 13, 920–921.
5. Ishikawa NK, Kasuya MCM and Vanetti MCD (2001). Antibacterial Activity of *Lentinula edodes* grown in liquid medium. Brazilian J. Microbio. 32:206-210.

6. Lewis SM, Mayhugh MA, Freni SC, Thorn B, Cardoso S, Buffington C *et al.* (2003). Assessment of antioxidant nutrient intake of a population of southern US African-American and Caucasian women of various ages when compared to dietary reference intakes. J Nutr Health Aging 7, 121–128.
7. Lakshmi B, Tilak, JC, Adhikari S, Devasagayam TPA and Janardhanan KK (2005) Inhibition of lipid peroxidation induced by γ- radiation and AAPH in rat liver and brain mitochondria by mushrooms. Research communications Current Science Vol, 88, No. 3. 484-488.
8. Taira K, Miyashita Y, Okamoto K., Arimoto, S, Takahashi E and Negishi T. (2005), Novel anti mutagenic factors derived from the edible mushroom *Agrocybe cylindracea*, Mutation Research, Genetic Toxicology and Environmental Mutagenesis, 586(2):115-23.
9. Mattam JG., Ashiq M, Stantley J and Prema K (2007) Micronutrient composition of Edible Mushrooms, Wide Spectrum, 17-19.
10. Okhawa H, Ohishi N and Yagi K (1979) Assay for lipoperoxides in animal tissues by thiobarbituric acid reaction, Anal, Biochem, 95, 351-358.
11. Dodge JT, Mitchel C and Hanghan V (1963) The preparation and chemical characteristics of haemoglobin free erythrocytes, Arch. Biochem. Biophysics, 100, 119-130.
12. Brand-Williams W, Cuvelier M and Berset C (1999) Use of a free radical method to evaluate antioxidant activity, Lebensmitt-Wissenschaft und Technolog 28, 25-30.
13. Pulido R, Bravo, L and Saara- Calixto, F (2000) Antioxidant activity of Dietary polyphenols as determined by a modified Ferric Reducing / Antioxidant power assay, J. Agric. Food. Chem, 48, 3396-3402.
14. Turkoglu A, Kivak I, Mercan N, Duru ME, Gezer K and Turkoglu H (2006), Antioxidant and Antimicrobial activities of *Morchella conica* Pers, African Journal of Biotechnology vol 5 (11), 1146-1150.
15. Halliwell B & Gutteridge J (1981) Formation of a thiobarbituric-acid-reactive substance from deoxyribose in the presence of iron salts. The role of superoxide and hydroxyl radicals. FEBS Letters, 128: 347-352.
16. Fraga C, Leibovitz B & Tappel, A (1987) Halogenated compounds as inducers of lipid peroxidation in tissue slices, Free Radical Biology and Medicine 3, 119-123.
17. Khanam S, Shivprasad HN and Kshama D (2004) Invitro antioxidant screening models: a review, Indian J Pharm Educ, 38, 180.
18. Khajuria A (1997) Lipid peroxidation, Everyman's Sci., Vol. XXXII, 109-113.
