



RESEARCH ARTICLE

CAST PARTIAL DENTURE USING PRECISION ATTACHMENT: A CASE REPORT

Rutuja Chavan¹ Abhishek Darekar² Susheel Chauthmal³ Jyotsna Deore⁴ and Akshay Mehtre⁵

¹PG Resident, SMBT Institute of Dental Sciences and Research, Igatpuri

²Professor & HOD, SMBT Institute of Dental Sciences and Research, Igatpuri

^{3,4,5}PG Resident, SMBT Institute of Dental Sciences and Research, Igatpuri

ARTICLE INFO

Article History:

Received 09th February, 2025

Received in revised form

21st March, 2025

Accepted 19th April, 2025

Published online 30th May, 2025

Key words:

Benign Jaw Tumor, Cemento-Ossifying Fibroma, Oral Soft Tissue Lesion, fibro-Osseous Growth, Non-Odontogenic Tumor.

*Corresponding author:

Rutuja Chavan

ABSTRACT

Aim: The aim of this case report is to demonstrate the clinical and prosthodontic management of a partially edentulous patient using a cast partial denture with precision attachments, emphasizing its advantages in improving retention, esthetics, and patient comfort compared to conventional clasp-retained removable partial dentures. **Methods:** This clinical report shows how to use the extra coronal attachment, precisagix, to restore a partially edentulous mandibular arch with detachable partial dentures. A matching fitting or housing was integrated into the removable prosthesis along with the crown, which allowed for some movement between the two parts of the prosthesis. This served as a non-rigid stress breaker and assisted in distributing the occlusal load. **Conclusion:** Semi-precision attachment retained cast partial dentures would be a great choice in situations when aesthetics are a top priority and the financial situation precludes the use of dental implants. The success of a distal extension cast partial denture depends on the abutment's ability to regulate stress, which is accomplished by using precise impression technique, wide coverage, a sturdy denture base, abutment splinting, and appropriate attachment selection. Semi-precision attachments are a feasible option that can enhance function, retention, and appearance with appropriate case selection and treatment planning.

Copyright©2025, Rutuja Chavan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Rutuja Chavan, Abhishek Darekar, Susheel Chauthmal, Jyotsna Deore and Akshay Mehtre. 2025. "Cast Partial Denture using Precision Attachment: A Case Report. International Journal of Current Research, 17, (05), 33024-33031.

INTRODUCTION

A retainer made up of a metal container and a closely fitting component is called an attachment. The male "patix" component is connected to a pontic or the denture framework, while the female "matrix" component is typically contained inside the extended or normal contours of the abutment tooth's crown. They are separated into two categories: semiprecision and precision attachment. All precision attachments have a very accurate smooth movement and are prefabricated, machined to extremely fine tolerances. They are often adjustable, some contain elements that can be replaced, and some have many functions like switching from friction retention to snap retention.^{1,2} Furthermore, some are resistive to a certain level of unfavourable stress transfer to the abutment tooth. These attachment styles are referred to as "stress directors," Rotatory movement (movement in numerous planes) or hinge-like movement (movement in one plane) are the two types of flexible attachments. Semi-precision attachments are formed using patterns made of wax, nylon, or plastic, or they can be hand waxed by a dental laboratory worker. Because of the nature of their creation, they have less exact tolerances.^{3,4} Attachments are further divided into two categories. Intra-coronal refers to a semi-precision or precision attachment that is situated inside the abutment tooth's

contours, whereas extra-coronal refers to an attachment that is situated outside of them.⁵ This clinical report shows how to use the extra coronal attachment, precisagix, to restore a partially edentulous mandibular arch with detachable partial dentures. A matching fitting or housing was integrated into the removable prosthesis along with the crown, which allowed for some movement between the two parts of the prosthesis. This served as a non-rigid stress breaker and assisted in distributing the occlusal load.

CASE REPORT

A 53 year old male patient reported to the Department of Prosthodontics and Crown and Bridge with complaint of missing teeth. Extraoral examination revealed facial asymmetry. On intraoral examination teeth numbers #14, 24, 25, 26, 27, 46, were found missing, there was generalised attrition, silver amalgam filling with 16, facets with upper and lower posterior teeth were found missing. Past dental history revealed, Patient had undergone multiple extraction in maxillary and mandibular region. After clinical and radiological examination and explaining the pros and cons of the treatment options it was decided to rehabilitate the case with attachment retained cast partial denture. To create the

diagnostic cast, an irreversible hydrocolloid imprint material was used to take a primary impression, which was then poured into a dental stone. Ceka Sagix attachment placement was intended for distal surface of 23 And a 1.7 mm spherical male was planned to be placed. The first master cast was created by preparing tooth number 23 by taking an impression using extra silicone impression material and pouring it into a die stone. The plastic male portion of the attachment was affixed to the mesial side of the wax patterns after the prepared teeth were waxed up in the standard way. The correct insertion path must be chosen in order to enter the cast plastic male.



Pre Operative A] Maxillary Arch



Pre Operative; A] Mandibular Arch



Pre operative intraoral view



Occlusion on right side



Occlusion on left side



Fig 1. Diagnostic surveying



Fig 2. JCEKA PRECI SAGIX attachment

Following the attachment of a paralleling mandrel to the surveyor, the plastic males were integrated into the prepared tooth's wax-up. All of the males' paths were drawn parallel to one another. When using detachable partial dentures, a lingual bracing arm is advised. A hard dental alloy was used for the casting process. An impression of the male portion was made at the subsequent visit using extra silicone impression material. The male part's impression was placed in cold-cured acrylic resin, and die stone was used for the remaining portion. This made sure the male portion wouldn't shatter during the retrieval of the second master cast. The female is being



Fig. 4. Tooth preparation with 23 for metal ceramic crown



Fig. 5. Elastomeric impression



Fig. 6. Wax pattern for coping along with preci sagix plastic male attachment

processed. Lay the pink replicating dummies on top of the males and line them up in a line. Remove the duplication in the refractory material and the undercuts. Complete the cast frame without causing any harm to the female's cavity's retentive ledge. The access opening should be highly polished. A framework try-in was conducted. A record of the jaw connection and a tooth arrangement were established. Once the denture has been tried on, process the acrylic resin by pressing the pink processing female into the frame with the insertion tool.




Fig. 7. Spruing of wax pattern



Fig. 8. Spruing in the investment ring




Fig. 9. Casted coping along with preci sagix male attachment

Changing the processing female: To swap out a processing female, remove it using a pointed tool. Insert a fresh female using the insertion tool. There are three retention levels available:



Fig. 10. Temporary crown cementation and rest seat preparation on 13, 16



Fig. 11. Pick up impression



Fig. 12. Retrieved cast



Fig. 13. Refractory cast



Fig. 14. Wax pattern fabrication for cast partial framework



Fig. 15. Spruing the wax pattern



Fig. 16. Investing the pattern



Fig. 17. Retrived casting

Yellow: typical retention **White:** decreased recall **Red:** higher retention. The smile's postoperative appearance demonstrated a noticeable improvement. Recall visits were made at one week interval and regular oral hygiene instructions were explained to the patient.

DISCUSSION

Direct casting of plastic wax or a refractory pattern is used to create semi-precision attachments. It offers the remarkable



Fig. 18. After sprue cutting




Fig. 19. After finishing and polishing



Fig. 20. After border moulding in green stick impression compound



Fig. 21. Wash impression in light body elastomeric impression material



Fig. 22. Pick up impression




Fig. 23. Bite recorded on wax rim



Fig. 24. Try in



Fig. 25. Flasking procedure


**Fig. 26. Face bow Recorded****Fig 27| Acrylised cast partial denture****Fig. 28. Placement of female processing clip****Fig. 29. CPD Insertion**



Pre Operative



Post Operative



Post Operative

benefits of increased comfort, retention, and aesthetics as well as less postoperative cast partial denture corrections. Usually recommended for nonparallel abutments, distal extension bases, and long span edentulous arches. Precision attachments were developed in response to the need to balance functional stability and aesthetic appeal in partial dentures. The size and form of the abutment teeth should be taken into consideration when choosing between an intracoronal and an extracoronal attachment. Compared to extracoronal attachments, intracoronal attachments necessitate greater tooth reduction and preparation.⁵ Here, the extracoronal male and female components made up the preci-sagix. Castable plastic males that are integrated into the wax-up with paralleling mandrels make up Sagix. There was a 2.2 mm diameter sagix utilised.^{6,7} Three levels of retention are offered for female components: The colours red, white, and yellow indicate increased, decreased, and normal retention, respectively. The excessive tooth reduction, compromised embrasures, and poor aesthetics that occur with using intracoronal attachments are all addressed with semi-precision attachments. The other benefits include a large selection of alloys and simplicity of handling. Each resilient counterpart attachment has a shelf life specified by the makers. The plastic resilient cap needs to be replaced when its retentive capacity is depleted due to wear and tear from use.⁷⁻⁹

CONCLUSION

Semi-precision attachment retained cast partial dentures would be a great choice in situations when aesthetics are a top priority and the financial situation precludes the use of dental implants. The success of a distal extension cast partial denture depends on the abutment's ability to regulate stress, which is accomplished by using precise impression technique, wide coverage, a sturdy denture base, abutment splinting, and appropriate attachment selection. Semi-precision attachments are a feasible option that can enhance function, retention, and appearance with appropriate case selection and treatment planning.

REFERENCES

1. The glossary of prosthodontic terms. 9th edition. J Prosthet Dent 2017;117: e1-e105.
2. Burns DR, Ward T. A review of attachments for removable partial denture design: Part Classification and selection. Int J Prosthodont 1990;3:98-102.
3. Becerra G, MacEntee M. A classification of precision attachments. J Prosthet Dent 1987;58:322-327.
4. Preiskel HW. Precision Attachments in Prosthodontics. Vol 1: The applications of intracoronal and extracoronal attachments Chicago: Quintessence, 1984.
5. Gozneli R, Yildiz C, Vanlioglu B, Evren BA, Kulak-Ozkan Y. Retention behaviors of
6. different attachment systems: Precious versus nonprecious, precision versus semi-precision. Dent Mater J. 2013;32:801-7.
7. Thapa D, Shrestha P. Removable prosthesis with extra coronal attachment for the management of distal extension case: a case report. J Nep Prosthodont Soc (JNPS) 2018; 2:100-04.
8. Pandey B, Joshi SP, Thapa D, Shrestha P. Use of a semi-precision attachment to fabricate a removable partial

- denture: a case report. Journal of Nepalese Prosthodontic Society. 2019 Dec 31;2(2):108-12.
9. Hedzelek W, Rzatowski S, Czarnecka B. Evaluation of the retentive characteristics of semi-precision extracoronal attachments. J Oral Rehab 2011; 38:462-68.
10. Patil S, Kulkarni S, Thakur S, Naik B. Crown lengthening procedure following intentional endodontic therapy for correction of supraerupted posterior teeth: Case series with long-term follow-up. J Ind Soc Periodontol 2016;20:103-8.
