



ISSN: 0975-833X

Available online at <http://www.journalcra.com>

International Journal of Current Research
Vol.3, Issue, 4, pp.259-262, April, 2011

INTERNATIONAL JOURNAL
OF CURRENT RESEARCH

RESEARCH ARTICLE

QUANTITATIVE VARIATION OF PROTEIN IN THE TISSUES OF A FRESH WATER FISH *Clarias batrachus* EXPOSED TO MERCURY AND CHROMIUM

Mary Josephine Rani A¹, John Milton MC^{2*}, Uthiralingam M² and Azhaguraj R²

¹Department of Zoology, Auxilium College, Katpadi, Vellore 632006.

²PG and Research Department of Advanced Zoology and Biotechnology, Loyola College, Chennai 600 034, INDIA.

ARTICLE INFO

Article History:

Received 5th February, 2011
Received in revised form
24th February, 2011
Accepted 15th March, 2011
Published online 27th April 2011

Key Words:

Clarias batrachus,
Mercury,
Chromium, Protein, Bradford

ABSTRACT

The present study was carried out to determine the effects of sub-lethal concentration (96 h-LC₅₀) of mercury (0.026ppm, 0.040ppm and 0.080ppm) and chromium (3.4ppm, 5.1ppm and 10.2ppm) for 28 days in *Clarias batrachus* and the protein content in the brain, gills, liver, kidney and muscle were quantified Bradford method. The levels of protein content in the liver, kidney, brain, muscle and gill showed fluctuations of decreasing and increasing trend in fishes exposed to different concentrations of mercury and chromium in a time-dose dependent manner. The protein variation in the tissues of *C. batrachus* were significant when exposed in mercury and chromium. The variations observed in the total protein quantified in the *C. batrachus* as a general indicator of pollutant induced stress response is discussed.

© Copy Right, IJCR, 2011 Academic Journals. All rights reserved.

INTRODUCTION

The contamination of aquatic ecosystem with a wide range of pollutants has become a matter of great concern over the last few decades, not only because of the threat to public water supplies, but also with of the damage caused to the aquatic life. The river systems may be excessively contaminated with heavy metals released from domestic, industrial, mining and agricultural effluents (Canli and Kalay, 1998). Mercury (Hg) is a highly toxic, nonessential, persistent, immutable and non-biodegradable metal that undergoes many changes during transfer through different trophic levels of the food chain. Hg occurs naturally in the environment and exists in several forms such as metallic mercury, inorganic mercury and organic mercury. Chromium is one of the heavy metal pollutants produced from ferrochrome, tanning and pigment industries. Chromium in nature exists in toxic Cr+4 and Cr+3 forms. The Cr+6 ion is known to affect several plant processes. Cr+3 is readily oxidized to Cr+6 in soils because of the presence of oxidized Mn, which serves as the electron acceptor in the reaction. Heavy metals can be absorbed from the soil and atmosphere, accumulate in the organs of the plants and show their phytotoxic effects (Parmar and Chanda, 2005).

Clarias batrachus (Linn.) is an air-breathing catfish, *C. batrachus* occurs in fresh and brackish waters throughout India. It is highly valued as a table fish throughout the Indian subcontinent and is preferred for culture even in muddy and shallow waters (Debnath and Surajit, 2009). Protein plays a vital role in the physiology of living organisms. All biological activities are regulated by enzymes and hormones, which are also proteins. Therefore the assessment of the protein content and amino-acids of the various internal organs can be

considered as a diagnostic tool to determine the physiological phases of the cell (Kapila and Ragothaman, 1999). Dinodia *et al.* (2002) observed marked reduction in residual protein levels in *Labeo rohita*, *Cirrhinus mrigala* and *Cyprinus carpio* due to cadmium toxicity. Srivastava *et al.* (2002) recorded changes in protein biochemistry in the liver and muscle of *Channa punctatus* due to zinc toxicity. Protein content was significantly decreased in liver and muscle of *C. carpio* (Dhanapackiam *et al.*, 2000) and *C. batrachus* (Aruna *et al.*, 2000) under stress. Studies by Neha *et al.* (2004) on tallest *Labeo rohita* exposed to chromium chloride showed that there was continuous time dependent depletion in the protein content and maximum protein was depleted after the exposure to longer duration. However, in nickel chloride and zinc chloride treated fish, there was a decrease in protein content initially but after the longer exposure period; increase in tissue protein was noted. It has now been well established that heavy metals even in traces interfere with various physiological and metabolic processes.

MATERIALS AND METHODS

Healthy living specimens of *Clarias batrachus* were procured, from Bharath Fish Seed India, Poondi, Thiruvallur district, Tamilnadu. *C. batrachus* measuring 9.5± 0.5cm in length and weighing 5 ± 0.5 gm were used in the experiment. The fishes were acclimatized to the laboratory conditions for 20 days prior to experiment in a glass aquarium (40 x 40 x 100 cm) filled with dechlorinated water. Water quality characteristics were determined. The mean values for the test water qualities were as follows, temperature 27.5±1.5°C, pH 7.5±0.03, dissolved oxygen 6.4±0.2mg/l, alkalinity 250±2.8mg/l as CaCO₃, total hardness 456±3.5mg/l. Stock solutions of mercuric chloride and potassium dichromate were prepared by dissolving analytical grade mercuric chloride (HgCl₂ from Merck) and

*Corresponding author: djohnmilton@gmail.com

potassium dichromate (K₂Cr₂O₇·7H₂O from Merck) respectively in double distilled water. Acute toxicity test was conducted in accordance with standard methods (APHA, 1995). Sublethal or safe level concentrations were derived from 96h LC₅₀ as per the procedure described by Finney (1971). The 96 h-LC₅₀ (~ 10%) of were selected as sublethal concentrations of mercury (0.026ppm, 0.040ppm and 0.080ppm) and chromium (3.4ppm, 5.1ppm and 10.2ppm) and ten fishes of same size were exposed to each concentrations with three replicates for a period of 28 days. A control batch corresponding to each test group was simultaneously maintained. The experiments were repeated five times for each test concentrations. To maintain a constant toxic concentration in the media, the water medium with appropriate concentration of heavy metal was changed every day. Fish were fed with commercial pelleted diets *ad libitum*. After 7, 14, 21 and 28 days the gills, liver, kidney, brain and muscle tissues of control and treated *C. batrachus* were dissected and homogenized thoroughly with 5 ml of phosphate buffer (pH 7.8) and centrifuged at 10,000 rpm for 15 minutes. After centrifugation the supernatant was collected and the proteins were precipitated by adding equal volume of 10% ice cold TCA. The pellet was re-dissolved in 2 ml of 1N NaOH. To 0.1 ml of the above sample 5 ml of the protein reagent was added, mixed thoroughly and the absorbance was measured at 595 nm. Bovine serum albumin was used as standard (Bradford, 1976).The results were analyzed using SPSS 10.1 version ANOVA followed by post-hoc Tukey-HSD's test, the significant levels of difference between control and treated fish were calculated at p < 0.05 level.

are 120 ppm, 115 ppm, 110 ppm and 102 ppm respectively. Increase in exposure period results in decreased LC₅₀ values. The sublethal concentrations (96 h-LC₅₀) of mercury (0.026ppm, 0.040ppm and 0.080ppm) and 96 h-LC₅₀) chromium (3.4ppm, 5.1ppm and 10.2ppm) were treated to *C. batrachus* for 28 days. The highest amount of protein was observed in Liver, kidney, brain, muscle, and gills in the control group and compared with the mercury and chromium treated fishes and tabulated (Tables 1&2). Gills, kidney, brain, muscle and liver showed pronounced changes in the protein content as sublethal mercury and chromium concentration increased. As the duration of the exposure period increased, there was a sudden decrease from 7th day to the 14th day followed by slight increase from 14th to 21st and to 28th day at 0.026 ppm, 0.04ppm and 0.08ppm sublethal concentrations of mercury. All the obtained values were subjected to statistical analysis and the values were significant at p < 0.05 level. Results indicate that sublethal mercury and chromium exposure leads to a general decline and gradual increase in the level of proteins in liver, kidney, brain, muscle and gill tissues of *C. batrachus*.

DISCUSSION

Proteins are highly sensitive to heavy metals and are one of the earliest indicators of heavy metal poisoning (Kakkar and Jaffery, 2005). Heavy metals may alter the protein concentration through impairing the synthesis and metabolism

Table 1. The levels of protein content (µl/ml) in tissues of *Clarias batrachus* exposed to sublethal concentrations (% 96h LC₅₀) of mercury (ppm) toxicity (7 to 28 days).

Days Tissues	7 days				14 Days				21 Days				28 Days			
	Cont	0.026	0.040	0.080												
Gill	2.64 ±0.29	2.25 ±0.07	2.17 ±0.06	1.94 ±0.02	2.80 ±0.12	2.03 ±0.06	1.65 ±0.48	1.77 ±0.02	2.88 ±0.07	2.21 ±0.01	2.08 ±0.01	1.92 ±0.04	2.77 ±0.11	2.23 ±0.03	2.07 ±0.06	1.93 ±0.02
Kidney	4.18 ±0.01	3.17 ±0.03	2.86 ±0.03	2.47 ±0.02	4.18 ±0.01	2.82 ±0.01	2.59 ±0.01	1.97 ±0.12	4.13 ±0.01	3.06 ±0.01	2.61 ±0.01	2.27 ±0.02	4.19 ±0.01	3.10 ±0.01	2.86 ±0.07	2.31 ±0.02
Brain	3.88 ±0.02	2.47 ±0.19	2.2 ±0.05	2.06 ±0.02	3.82 ±0.05	2.07 ±0.02	2.09 ±0.01	1.90 ±0.01	3.78 ±0.09	2.18 ±0.02	2.11 ±0.01	2.10 ±0.01	3.87 ±0.06	2.23 ±0.03	2.14 ±0.11	2.18 ±0.07
Muscle	3.25 ±0.02	2.73 ±0.46	2.40 ±0.01	2.43 ±0.47	3.26 ±0.06	2.06 ±0.02	2.01 ±0.01	1.89 ±0.01	3.19 ±0.04	2.19 ±0.01	2.10 ±0.01	2.01 ±0.01	3.16 ±0.27	2.31 ±0.01	2.28 ±0.01	2.10 ±0.01
Liver	4.43 ±0.05	3.26 ±0.03	3.09 ±0.01	2.56 ±0.03	4.37 ±0.02	2.9 ±0.04	2.82 ±0.04	2.04 ±0.05	4.51 ±0.01	3.06 ±0.05	2.90 ±0.01	2.39 ±0.01	4.45 ±0.03	3.10 ±0.01	3.04 ±0.04	2.45 ±0.03

Cont.- Control

Table 2. The levels of protein content (µl/ml) in tissues of *Clarias batrachus* exposed to sublethal concentrations (% 96h LC₅₀) of chromium (ppm) toxicity (7 to 28 days)

Days Tissues	7 days				14 Days				21 Days				28 Days			
	Cont	3.4	5.1	10.2	Cont	3.4	5.1	10.2	Cont	3.4	5.1	10.2	Cont	3.4	5.1	10.2
Gill	3.04 ±0.20	2.77 ±0.02	2.57 ±0.05	2.44 ±0.05	2.87 ±0.18	2.4 ±0.01	2.20 ±0.001	1.91 ±0.01	2.95 ±0.30	2.60 ±0.01	2.42 ±0.01	2.03 ±0.01	2.96 ±0.28	2.70 ±0.01	2.50 ±0.01	1.97 ±0.02
Kidney	3.86 ±1.14	3.3 ±0.03	3.09 ±0.01	2.77 ±0.01	4.27 ±0.08	2.85 ±0.04	2.64 ±0.03	2.57 ±0.02	4.27 ±0.09	3.05 ±0.03	2.92 ±0.02	2.67 ±0.01	4.57 ±0.05	3.11 ±0.01	3.01 ±0.01	2.70 ±0.01
Brain	3.61 ±0.15	2.96 ±0.01	2.82 ±0.06	2.60 ±0.02	3.81 ±0.10	2.33 ±0.01	2.18 ±0.01	1.86 ±0.04	3.74 ±0.12	2.83 ±0.04	2.63 ±0.01	2.43 ±0.04	4.25 ±1.83	3.04 ±0.02	2.98 ±0.01	2.63 ±0.01
Muscle	3.47 ±0.22	2.96 ±0.01	2.82 ±0.06	2.60 ±0.01	3.50 ±0.26	2.48 ±0.01	2.17 ±0.03	1.93 ±0.02	3.43 ±0.23	2.64 ±0.03	2.46 ±0.03	2.23 ±0.05	3.48 ±0.18	2.83 ±0.01	2.70 ±0.01	2.12 ±0.03
Liver	4.61 ±0.07	3.47 ±0.01	3.23 ±0.03	2.98 ±0.02	4.67 ±0.08	2.94 ±0.04	2.74 ±0.04	2.67 ±0.03	4.51 ±0.02	2.83 ±0.59	3.03 ±0.02	2.76 ±0.02	4.76 ±0.08	3.27 ±0.02	3.12 ±0.01	2.91 ±0.01

Cont.- Control

RESULTS

The median lethal concentration (LC₅₀) of mercury to *Clarias batrachus* for 24, 48, 72 and 96 h of exposure were 1.4 ppm, 1.2 ppm, 1.0 ppm and 0.8 ppm respectively. The LC₅₀ value of chromium to *C. batrachus* for 24, 48, 72 and 96 h of exposure

of protein, DNA and RNA as well as alter the activity of lysosomal enzymes (Jana and Bandopadhyaya, 1987). It is possible that pollutant stress influences the conversion of tissue proteins into soluble fractions reaching the blood and hence, decrease in protein content was observed in the tissues of liver and muscle of *Labeo rohita* exposed chronically to heavy metal

salts, chromium chloride, nickel chloride and zinc chloride (Neha *et al.*, 2004). In the present study mercury and chromium sublethal toxicity stress also led to depletion of the original protein content in gill, kidney, brain, muscle and liver of *C. batrachus* was observed in the control and treated tissues (Table 1 & 2). Depletion of protein content has been observed in all the organs of *C. batrachus* till 14 days of exposure followed by slight increase from 14th to 21st and to 28th day after treatment. However, this increase could not reach the level of the control values. The maximum level of protein was found in the liver and the minimum in the gills and muscles of *C. batrachus*. Increase in exposure period as well as sublethal mercury concentration led to decrease in total protein content though their levels show replenishment after 21 days of exposure to the toxicant. Similar observation was noticed in the chromium treated tissues of the experimental fish.

Kapila and Ragothaman (1999) have also reported decreased tissue proteins followed by increased level of proteins in the fish, *Boleophthalmus dussumieri* exposed to mercury, copper and cadmium for prolonged periods. The initial drop in the protein content during mercury toxicity may be on account of reduced protein synthesis and an enhanced proteolysis in the various organs of fish (Jagadeesan and Mathivanan, 1999). The depletion of protein level induces diversification of energy to meet the impending energy demands during the toxic stress. The reduction in tissue proteins reflects a prior increased energy cost of homeostasis, tissue repair and detoxification under toxic stress. It is also possible that when an animal is under toxic stress, diversification of energy occurs to accomplish the impending energy demands. Hence depletion in protein level is observed (Neff, 1985). Evidently the whole energy is required to mitigate any stress condition and this energy may be derived from proteins (Shakoori *et al.*, 1992).

The obvious reasons for the varying proteins levels in fish under heavy metal toxicity could be due to the rapid metabolism under heavy metal stress (Shakoori *et al.*, 1992). Reduction in protein content is a cause of decrease in RNA in the tissues of fish or amino-acid incorporation and disaggregation of polysomes (Holbrook, 1980). The induced proteolysis by the heavy metal in the respective tissues contributes to the increase in the amino-acids (Radhaiah *et al.*, 1987; Ravichandran *et al.*, 1994). During protein metabolism the removal of the amino group from different amino-acids was observed suggesting the elevated levels of amino-acids (Begum, 2008). Both transamination and deamination seems to be enhanced at this stage. Baskaran (1991) reported depletion of protein content in the muscle and liver of *Oreochromis mossambicus*, *Mystus vittatus* and *C. striatus* exposed to fenvalerate. Jeba kumar *et al.* (1990) observed decrease in protein content of *Lepidocephalichthys thermalis* exposed to sublethal concentrations of fenvalerate. A significant decrease was reported in the protein content of almost all tissues in *Ctenopharyngodon idellus* when exposed to sublethal and lethal concentrations of fenvalerate (Tilak *et al.*, 2001). The decrease of the protein content in the liver may be due to necrosis of the hepatocytes. Kondal *et al.* (1988) attributed the decrease in protein content of the liver of *Saccobranchus fossilis* and *Heteropneustes fossilis* exposed to factory effluent and ammonia respectively, to the availability of energy required for the synthesis of proteins as a consequence of the decreased level of the key enzyme (Malate dehydrogenase) and due to

enhanced catabolism of amino-acids, as a consequence of increased activity of glutamate dehydrogenase. Protein content significantly decreased in liver and muscle of *Cyprinus carpio communis* under stress (Dhanapackiam *et al.*, 2000) and *C. batrachus* (Aruna *et al.*, 2000).

Initial increase of protein content after 7th day of treatment reflects an increase in total enzyme activity of the tissue in order to match the physiological demand of the system. The fish needs to balance the adverse stress by synthesizing certain molecules which appear to be proteinaceous in nature to reduce the toxicity. The increase of protein content in the kidney reflects stimulated protein synthesis of detoxification enzymes at the expense of glycogen to meet additional energy requirements in the synthetic activity of the tissues (Aruna, *et al.*, 2000). The kidney which is an important organ of excretion and osmoregulation is indirectly affected by pollution through blood circulation (Newman and MacLean, 1974). Decreased protein content in the kidney could be possible due to protein break down leading to increased amino-acid pool of tissue (Radhaiah *et al.*, 1987). The decrease in the protein content of the kidney of *C. batrachus* after exposure to malathion is attributed to the impairment of protein synthesis and/or increase in the rate of their degradation to amino acids which may be fed to the TCA cycle through aminotransferase probably to cope with the stress condition (Aruna *et al.*, 2000). Kasturi and Chandran (1997) observed decreased level of protein content in the tissues of *Mystus gulio* exposed to sublethal concentration of lead. Sen *et al.* (1992) also observed decrease in protein content of brain and liver of *C. punctatus* exposed to toxic effects of zinc. These observations are in accordance with the results of the present study.

Conclusion

In this study, mercury and chromium sub lethal concentrations in different tissues of *C. batrachus* showed different elevation of the protein content which shows the toxicological effects of heavy metals on fish that can be used for monitoring the heavy metal toxicity in aquatic ecosystem.

REFERENCES

- APHA, 1995. Standard Methods for the Examination of water and wastewater, 19th edition, American Public Health Association, Washington DC.
- Aruna, K., Singh, S. and Shrivastava, K. 2000. Malathion induced biochemical changes in the kidney of freshwater fish *Clarias batrachus*. *J. Ecotoxicol & Environ.*, 10(1): 11-14.
- Baskaran, P. 1991. Use of biochemical parameters in biomonitoring of pesticide pollution in some freshwater fishes. *J. Ecotoxicol. Environ. Monit.*, 1(2): 101-104.
- Begum, G. 2008. Assessment of biochemical markers of Carbofuran toxicity and recovery response in tissues of the freshwater teleost, *Clarias batrachus* (Linn). *Bull Environ. Contam. Toxicol.* 81:480-484.
- Bradford, M. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein binding. *Anal. Biochem.*, 72:248 - 254.
- Canli, M.O., and Kalay, M. 1998. Levels of Heavy Metals (Cd, Pb, Cu, Cr and Ni) in Tissue of *Cyprinus carpio*, *Barbus*

- capito* and *Chondrostoma regium* from the Seyhan River, Turkey. *Tr. J. of Zoology*, 22 : 149-157
- Debnath, and Surajit, 2009 .Traditional Consumption of Magur (*Clarias batrachus*) an Air Breathing Catfish among the Population of North Eastern India is Rationalized by Its Blood Lipid Parameters (May 9, 2009). CHEMREFERENCE, Annual Research Symposium, IIT Madras 22 - 23rd August, 2009. Available at SSRN:<http://ssrn.com/abstract=1401145>.
- Dhanapackiam, P., Fathima Bibi, N. and Ramasamy, V.K. 2000. Effect of malathion on a freshwater fish *Cyprinus carpio*. *Uttar Pradesh J. Zool.*, 20(1): 13-16.
- Dinodia, G.S., Jain, K.L. and Gupta, R.K. 2002. Effect of cadmium toxicity on total body carbohydrates and residual protein contents in some freshwater fishes *Himalayan J. Env. Zool.*, 16(2): 171-176.
- Finney, D. J. 1971. Probit analysis. 3d ed., Cambridge University Press, London and New York.
- Holbrook, Jr.D.J. 1980. Effects of toxicants on nucleic acid and protein metabolism In: Introduction to biochemical toxicology (eds.) Hodgson, E. and Guthrie, F.E. Blackwell Scientific Publications, Oxford, p.261 -284.
- Jagadeesan, G. and Mathivanan, A. 1999. Organic constituent's changes induced by three different sublethal concentrations of mercury and recovery in the liver tissue of *Labeo Rohita* fingerlings. *Poll.Res*. 18 (2): 177-181.
- Jana, S. and Bandopadhyaya, N. 1987. Effect of heavy metals on some biochemical parameters in the freshwater fish, *Channa punctatus*. *Environ. Ecol.*, 5(3): 488-493.
- Jayantharao, and Murthy, V.S.R. 1983. Hypoxia induced the haemopoietic tissue in freshwater fish. *Oreochromis mossambicus* exposed to phosphamidon. *Geobios*, 10: 264-267.
- Jeba Kumar, S.R.D., Flora, S.D.J., Ganessan, R. M., Jagadeesan, G. and Jaya Raman, J. 1990. Effect of short term sublethal exposure cypermethrin on the organic constituents of the freshwater fish, *Lepidocephalichthys thermails*. *J. Environ. Biol.*, 11(2): 203-209.
- Kakkar P. and Jaffery FN. 2005. Biological markers for metal toxicity. *Environ Toxicol. and Pharm.* 19: (2) 335-349.
- Kapila, M. and Regothaman, G. 1999. Mercury, copper and cadmium induced changes in the total protein level in muscle tissue of an edible estuarine, fish, *Belophthalmus dussumieri* (Cuv). *J. environ. Biol.*, 20(3): 231-234.
- Kasturi, J. and Chandran, M.R. 1997. Sublethal effect of lead on feeding, energetics, growth performance, biochemical composition and accumulation of the estuarine catfish, *Mystus gulio* (Hamilton). *J. Environ. Bio.*, 18(1): 95-101.
- Kondal, J.K., Saxena, P.K., and Soni, G.L. 1988. Effect of vegetable oil factory effluent on the liver of freshwater teleost, *Heteropneustes fossilis* (Bl.)-A biochemical study. *Biological wastes*, 25: 177-185.
- Neff, J.M. 1985. Use of biochemical measurement to detect pollutant mediated damage to fish. *ASTM Spec. Tech. Publ.*, 854: 155-183.
- Neha, B., Vankhede, G.N. and Dhande, R. R. 2004. Heavy metals induced biochemical alterations in freshwater fish *Labeo rohita*. *J. Ecotoxicology and Environmental monitoring*, 1: 14-15.
- Newman, M.W. and MacLean, 1974. Physiological response of the cunner *Tautoglabrus adspersus* to cadmium. VI Histopathology No. A Tech. Report NMFS, SSRF: p.681.
- Parmar.N.G and V. Chanda. 2005. Effects of Mercury and Chromium on Peroxidase and IAA Oxidase Enzymes in the Seedlings of *Phaseolus vulgaris*. *Turk J Biol.* 29: 15-21.
- Radhaiah, V., Girija, M. and Jayant Rao, K. 1987. Changes in selected biochemical parameters in the kidney and blood of the fish, *Tilapia mossambicus* exposed to heptachlore. *Bull. Environ. Contam. Toxicol.*, 39: 1006-1011.
- Ravichandran, S., Midhunashanthi, K. and Indira, N. 1994. Impact of phenol on protein metabolism in the freshwater fish *Oreochromis mossambicus*. *J. Ecotoxicol. Environ.*, 4(1): 33-37.
- Sen G., Behera, M.K., and Patel, P.N. 1992. Toxicity of zinc on liver and brain of the Fish *Channa punctatus* (Bloch). *Environ. Ecol.*, (10)3: 742-744.
- Shakoori, A.R., Ilyas, M. and Aziz, F. 1992. Toxicity of sublethal doses of Trebon (ethofenprox) on total blood serum proteins, acetylcholinesterase activity and SDS-PAGE pattern of blood serum proteins of *Cirrhinus mrigala*. *Pak. J. Zool.*, 24(3): 235-241.
- Srivastava, N., Kaushik, N. and Gupta, P. 2002. Zinc induced changes in the liver and muscle of fish *Channa punctatus* (Bloch). *J. Ecophysio. Occupl. Hlth.*, 2(3 & 4): 197-204.
- Tilak, K.S., Veeraiah, K. and Ramanakumari, G.V. 2001. Toxicity and effect of chloropyrifos to the fresh water fish, *Labeo rohita* (Hamilton). *J. Poll. Res.*, 20(3): 443 – 445.
