



ISSN: 0975-833X

RESEARCH ARTICLE

HETEROSIS AND COMBINING ABILITY ANALYSIS IN TETRAPLOID COTTON (*G. HIRSUTUM* AND *G. BARBADENSE* L.)

*¹Kannan, N. and ²Saravanan, K.

¹Principal Breeder (Cotton), Rasi seeds Pvt Ltd., Attur, India

²Department of Genetics and Plant Breeding, Annamalai University, Annamalai Nagar-608 002, India

ARTICLE INFO

Article History:

Received 10th February, 2015

Received in revised form

18th March, 2015

Accepted 24th April, 2015

Published online 31st May, 2015

Key words:

Cotton,
Lx T analysis,
Heterosis,
Combining ability,
Gene action.

ABSTRACT

The present investigation was carried out with a view to study the heterosis, combining ability and gene action for seed cotton yield, yield attributing traits, ginning outturn and important fibre quality parameters in tetraploid cotton (*G.hirsutum* L. and *G.barbadense* L.). A set of 31 genotypes consisting of 21 (HxB) hybrids, Seven Lines of *G. hirsutum* and three testers of *G. barbadense* and one check hybrid DCH 32 were grown at Rasi seeds Research and Development Farm Attur. The hybrids were developed adopting Lines and Testers mating design. Analysis of variance showed significant differences amongst parents and hybrids for all the characters indicating presence of genetic variability. Combining ability analysis indicated that both additive and non additive gene effects were important in the inheritance of all the traits. The ratio of variance due to GCA to that of SCA was less than one for all the character under study indicating importance of dominance gene effects in the inheritance of these characters. The parents CG 64 (*hirsutum*), CG67 (*hirsutum*) and CG45SB (*barbadense*) were good general combiner for the seed cotton yield per plant and number of bolls per plant. CG 45SB was also a good general combiner for the 2.5 per cent span length. The cross CG 64 x CG 45 SB recorded the highest *per se* performance and standard heterosis for seed cotton yield. Two crosses and CG64 x CG45SB and CG67 x CG45SB registered significant *per se* performance, positive *sca* effects along with significant positive standard heterosis for seed cotton yield and majority of yield components and fibre quality traits.

Copyright © 2015 Kannan, N. and Saravanan, K. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Kannan, N. and Saravanan, K. 2015. "Heterosis and combining ability analysis in Tetraploid cotton (*G. hirsutum* and *G. barbadense* L.)", *International Journal of Current Research*, 7, (5), 16590-16595.

INTRODUCTION

Cotton, the king of fibre is one of the most momentous and important cash crops of the country. It is the most important commercial crop contributing nearly 65 per cent of the total raw material needs of the textile industry in our country. Cotton is cultivated in area of 115.53 lakh hectares with the production of 375 lakh bales (AICCIP, Report, 2014). Among the four cultivated species of cotton the varieties and hybrids of two species *G.hirsutum*, *G.barbadense*, *hirsutum X hirsutum* and *hirsutum X barbadense* hybrids were widely cultivated in India. In India around 7 – 9 lakh bales of ELS cotton is needed for the textile mills but the production is only around 5.0 lakh bales. The inter specific hybrids (*G. hirsutum X G. barbadense*) which are the one of the source for ELS cotton production is to exploited to meet the demand. Several workers reported higher heterosis in case of inter-specific hybrids. Exploitation of heterosis (*G. hirsutum X G. barbadense*) on commercial scale and systemic varietal improvement through

hybridization are the main tools to increase ELS cotton production. Study of combining ability is important for selecting parents for hybridization. Sprague and Tatum (1942), first time proposed the concepts of general combining ability (*gca*) and specific combining ability (*sca*). According to them, GCA variance is due to additive variance and SCA variance is due to non-additive variance, both acts as an important diagnostic tool in selection of suitable parents and cross combination. Among the various design used for combining ability analysis, Line x Tester analysis (Kempthorne, 1957) has been extensively used to assess the combining ability of parents and crosses for different quantitative characters as well as to study the extent of heterosis for yield, yield contributing characters and fibre quality traits in cotton.

MATERIALS AND METHODS

The present investigation was carried out at Research and Development Farm of Rasi Seeds, Attur, during 2009 Winter. The experimental materials consisted of 21 hybrids (*hirsutum X barbadense*) obtained by crossing of seven lines (*hirsutum*) viz., CG62, CG64, CG67, CG91, CG92, CG150 and CG163 and

*Corresponding author: Kannan, N.

Principal Breeder (Cotton), Rasi seeds Pvt Ltd., Attur, India.

three testers (*barbadense*) viz., CG45SB, CG45E and CG305. The hybrids and parents were evaluated in randomized block design with two replications. Each treatment was raised four rows of 6.8m length spaced at 1.2m apart with plant to plant distance of 75cm. All the recommended agronomical practices and plant protection measures were followed as and when required to raise a good crop of cotton. Observations were recorded for number of bolls per plant, boll weight (g), seed cotton yield per plant (g), number of seeds per boll, ginning outturn (per cent), Lint Index (g), Seed Index (g), 2.5 Span Length (mm), Uniformity Ratio (per cent), Fibre Strength (3.2 g/tex), Micronaire and Elongation. Data were recorded on five random competitive plants from each entry from all replications and mean of five plants was taken for further analysis. Observations were recorded on number of bolls per plant, boll weight (g), seed cotton yield per plant (g), number of seeds per boll, ginning outturn (per cent), lint Index (g), seed Index (g), 2.5 per cent Span Length (mm), uniformity Ratio (per cent), fibre Strength (g/tex), Micronaire (10^{-6} g/inch) and elongation. Data were recorded on five random competitive plants from each entry from all replications and mean of five plants was taken for further analysis. Combining ability analysis were estimated as per Kempthorne (1957). Standard heterosis were estimated as per the procedure suggested by Shull (1948) and Liang *et al.* (1971).

RESULTS AND DISCUSSION

Analysis of variance showed highly significant differences due to genotypes for all the traits indicating the presence of sufficient variability in the experimental materials (Table 1). Parents and hybrids showed significant differences between all the characters studied. Significance of variance in parents various hybrids interaction provides adequacy for comparing the heterotic expression for all the characters except lint index. The ratio of variance due to GCA to that of SCA was lesser than one for all the characters which shows the predominance of dominant gene action for all the characters under study indicating the exploitation of heterosis by hybrid development is most suitable method of breeding. These findings are in agreement with those obtained by Simon *et al.* (2013).

Information on the *per se* performance and nature of general combining ability effects (*gca*) of characters is necessary for selection of suitable parents for developing hybrids. The *gca* effects of the parents along with their mean performance for various characters are given in Table 2 and 3. Based on the *per se* performance the *hirsutum* line CG163 (217.6g), was superior for the Seed cotton yield per plant followed by CG91 (217.15 g) and CG150 (207.9 g). The *barbadense* genotype CG45E was superior for 2.5 Span Length (41.10mm) and CG45 SB was superior for Fibre strength (36.9 g/tex). The parents CG 45SB (37.87 g), CG 64(31.94 g) and CG 67 (29.19 g) were good general combiners for the seed cotton yield per plant. These parents also recorded significant *gca* effects of 25.48, 24.90 and 19.57 for number of bolls per plant respectively. The genotype CG150 recorded highest significant *gca* effect (0.24) for boll weight, number of seeds per boll (1.45), ginning outturn (2.96) and lint index (1.26). For the character 2.5 per cent span length the genotype CG 45SB and CG 62 recorded highest significant *gca* effects (1.03).

The genotype CG 67 registered highest significant *gca* effect (2.63) followed by CG 62(2.30) and CG 45SB(1.41) for fibre strength. These results are in agreement with the findings of Rao and Gopinath (2013) and Amir *et al.*, (2012). The *sca* effects of the hybrids along with their mean performance for various characters are given in Table 4. The specific combining ability (*sca*) effect alone may not be the appropriate choice for exploitation of heterosis because the hybrid with low mean value may also possess high *sca* effect. Hence, the cross combinations were to be identified based on two criterias viz., *per se* performance and the gene action involved in the crosses for further exploitation. In the present investigation based on the *per se* performance the cross combination CG64 x CG 45 SB was found to be superior for seed cotton yield (353.4 g), number of bolls per plant (166.5), boll weight (4.4 g) and number of seeds per boll (25.5). This cross combination CG 64 x CG 45 SB recorded highest significant *sca* effects for boll weight (0.45), number of seeds per boll (3.31) and elongation (0.66). The hybrid combination CG150 x CG45E was superior for ginning outturn which recorded 33.55 per cent of mean performance of ginning outturn. For the character Lint index CG 150 x CG 45 E was superior for the *per se* performance (6.9 g) and *sca* effect (0.41). Regarding the fibre quality traits the cross combination CG 62 x CG 45SB for 2.5per cent span length (39.8mm) followed by CG 64 x CG 45SB (39.45mm). For the fibre strength the hybrid CG 64xCG 45SB was superior for mean performance (35.7 g/tex) and significant *sca* effect (1.72). Similar results were earlier reported by Vineela *et al.* (2012) and Amir *et al.* (2012).

A summarized account of the best parent *per se*, best general combiner, best F_1 *per se*, most heterotic crosses and best specific combination for various characters studied in the present investigation are presented in Table 5. The best performing cross CG 64 x CG 45SB recorded highest *per se* performance for seed cotton yield per plant (353.40 g) and Standard heterosis of 49.81 per cent, number of bolls per plant (166.50) and standard heterosis of 129.66 per cent, number of seeds per boll (25.50) and standard heterosis 2.00 per cent, uniformity ratio (47.15) and standard heterosis 9.65 per cent followed by the CG 67 x CG 45 SB for seed cotton yield, number of bolls per plant and fibre quality traits. All the parents involved in these crosses are having good general combining ability for seed cotton yield per plant and number of bolls per plant. The heterosis in this cross might have resulted from interaction of dominant gene contributed by both good combining parents. Pathak and Kumar (1975) reported close relationship between *gca* effects of parents and *sca* effect of their resultant crosses. In case of fibre quality analysis the cross CG 64 x CG 45 SB recorded 39.45 mm of 2.5per cent span length, and 33.15 g/tex of fibre strength and standard heterosis of 8.45 and 21.21 per cent respectively with the positive significant *sca* effects. The respective parents are also having significant positive *gca* effects for the 2.5 per cent span length and fibre strength. The next overall best hybrid CG 67 x CG 45 SB recorded high *per se* performance and standard heterosis for 2.5 per cent span length, fibre strength but negative *sca* effect for 2.5 per cent span length. Both the parents involved in this cross were having good general combining ability.

Table 1. Analysis of variance for combining ability for twelve characters in cotton

Source of variation	DF	Number of bolls per plant	Boll Weight (g)	Seed cotton yield per plant (g)	Number of seeds per boll	Ginning outturn (per cent)	Lint Index (g)	Seed Index (g)	2.5 Span Length (mm)	Uniformity Ratio (per cent)	Fibre Strength (g/tex)	Micronaire	Elongation
Cross	20	1542.18**	0.12**	3037.92**	6.50**	7.23**	1.01**	1.29**	3.53**	3.04**	17.27**	0.22**	0.62**
Line	6	2222.21**	0.14**	3701.64**	4.10**	17.83**	2.52**	1.91**	4.96**	3.73**	39.92**	0.27**	1.02**
Tester	2	7249.45**	0.18**	16411.07**	6.45**	0.77**	1.27**	2.24**	13.94**	10.47**	33.85**	0.52**	0.81**
LXT	12	250.95**	0.11**	477.20**	7.70**	3.01**	0.21**	0.83**	1.08**	1.46**	3.18**	0.15**	0.39**
Cross VS parents	1	21672.26**	5.17**	146733.33**	44.83**	116.65**	0.00	28.73**	97.00**	4.12**	72.40**	4.73**	0.08**
Error	30	13.74	0.01	5.69	0.78	0.05	0.01	0.02**	0.03	0.03	0.02	0.01	0.02
Var of GCA		50.44	0.00	100.03	0.05	0.16	0.03	0.02	0.10	0.06	0.55	0.00	0.01
Var of SCA		124.46	0.05	237.13	3.49	1.50	0.10	0.41	0.53	0.72	1.58	0.07	0.19
GCA/SCA		0.41	0.01	0.42	0.01	0.11	0.31	0.04	0.18	0.09	0.35	0.04	0.05

* Significant at 5 % probability level, ** Significant at 1 % probability level.

Table 2. Mean Performance of Lines and testers for twelve characters in cotton

PARENTS	Number of bolls per plant	Boll Weight (g)	Seed cotton yield per plant (g)	Number of seeds per boll	Ginning outturn (per cent)	Lint Index (g)	Seed Index (g)	2.5 Span Length (mm)	Uniformity Ratio (per cent)	Fibre Strength (g/tex)	Micronaire	Elongation
LINES												
CG64	65.5	4.55	157	26*	35**	5.65	10.2	30.8	46.25**	24.25	4.35**	5.85**
CG91	102**	4.65	217.15**	26.5*	38.2**	7**	11.65**	29.15	46.15**	22.05	4.65**	5.55*
CG150	90.5**	5.3**	207.9**	27.5**	42.4**	8.2**	12**	29.3	47.35**	24.05	4.85**	6.85**
CG62	66	5.35**	170.4	28.5**	32.25	5.15	10.75	36.05**	44.85	27	4.05**	5.35
CG67	72.5	5**	175.5**	26.5*	33.4*	5.35	10.65	36.3**	44.8	26.15	4.1**	5.05
CG92	66	5.45**	166.9	27*	35.65**	5.15	9.75	35.15*	45.05	26.1	3.45	4.65
CG163	76	5.65**	217.6**	30**	33*	5.15	9.95	33.55	42.65	25.75	3.4	5.55
TESTERS												
CG45SB	59.5	3.6	109.95	18	25.55	4.45	12.85**	40.5**	44.55	36.9**	2.8	4.55
CG45E	51.5	3.45	138.5	17.5	24.65	4.45	13.55**	41.1**	44.55	34.85**	3.15	4.4
CG305	46.5	3.75	104.65	19.5	28.6	5.15	12.35**	37.7	43.55	32.45**	3.25	5.3
Mean of Parents	69.60	4.67	166.55	24.70	32.87	5.57	11.37	34.96	44.98	27.96	3.81	5.31
CD 5%	7.56	0.19	4.86	1.79	0.43	0.21	0.28	0.35	0.32	0.29	0.18	0.26
CD 1%	10.19	0.26	6.56	2.42	0.59	0.28	0.38	0.47	0.43	0.39	0.24	0.35

* Significant at 5 % probability level, ** Significant at 1 % probability level.

Table 3. Estimates of general combining ability (gca) effects of Lines and testers for twelve characters in cotton

PARENTS	Number of bolls per plant	Boll Weight (g)	Seed cotton yield per plant (g)	Number of seeds per boll	Ginning outturn (per cent)	Lint Index (g)	Seed Index (g)	2.5 Span Length (mm)	Uniformity Ratio (%)	Fibre Strength (g/tex)	Micronaire	Elongation
LINES												
CG64	24.90 **	-0.22 **	31.94 **	-1.38 **	-0.02 ns	-0.18 **	0.12 **	0.16 *	1.08 **	1.05 **	0.05 ns	0.06 ns
CG91	-3.10 **	0.08 ns	0.38 ns	-0.21 ns	1.15 **	0.42 **	0.62 **	-1.30 **	-0.34 **	-3.48 **	0.14 **	0.31 **
CG150	-23.60 **	0.24 **	-28.02 **	1.45 **	2.96 **	1.26 **	0.37 **	-1.15 **	-1.31 **	-3.45 **	0.24 **	0.60 **
CG62	12.90 **	0.06 ns	7.16 **	0.12 ns	-1.92 **	-0.23 **	0.32 **	1.03 **	-0.36 **	2.30 **	-0.23 **	-0.35 **
CG67	19.57 **	-0.07 ns	29.19 **	0.12 ns	-2.00 **	-0.56 **	-0.14 **	0.88 **	0.33 **	2.63 **	-0.31 **	-0.65 **
CG92	-18.10 **	-0.11 *	-29.44 **	-0.05 ns	0.01 ns	-0.59 **	-1.09 **	0.18 *	-0.11 **	1.50 **	-0.06 ns	0.10 *
CG163	-12.60 **	0.03 ns	-11.21 **	-0.05 ns	-0.19 **	-0.13 **	-0.21 **	0.20 *	0.71 **	-0.55 **	0.19 **	-0.07 ns
TESTERS												
CG45SB	25.48 **	0.11 **	37.87 **	0.69 **	0.08 *	0.11 **	0.41 **	1.03 **	0.81 **	1.41 **	-0.11 **	-0.06 *
CG45E	-7.17 **	0.00 ns	-9.10 **	-0.02 ns	-0.26 **	-0.34 **	-0.39 **	-0.06 ns	0.10 **	0.25 **	-0.11 **	-0.21 **
CG305	-18.31 **	-0.11 **	-28.77 **	-0.67 **	0.19 **	0.23 **	-0.02 ns	-0.96 **	-0.91 **	-1.67 **	0.22 **	0.26 **

* Significant at 5 % probability level, ** Significant at 1 % probability level.

Table 4. *Per se* performance and estimates of specific combining ability (*sca*) effects of hybrids for twelve characters in cotton

	Number of bolls per plant		Boll weight (g)		Seed cotton yield per plant (g)		Numbr of seed per boll		Ginning outturn (per cent)		Lint intex (g)	
	MEAN	<i>sca</i>	MEAN	<i>sca</i>	MEAN	<i>sca</i>	MEAN	<i>sca</i>	MEAN	<i>sca</i>	MEAN	<i>sca</i>
CG64 X CG45SB	166.5**	6.52 **	4.4**	0.45 **	353.4**	12.96 **	25.5**	3.31 **	31.2**	1.20 **	5.45	-0.06 ns
CG64 X CG45E	141**	13.67 **	3.45	-0.38 **	308.55**	15.09 **	19.5	-1.98 **	28.35	-1.30 **	4.95	-0.11 ns
CG64 X CG305	96	-20.19 **	3.65	-0.07 ns	245.75	-28.05 **	19.5	-1.33 *	30.2*	0.10 ns	5.8*	0.17 *
CG91 X CG45SB	123.5**	-8.48 **	4.15	-0.10 ns	298.45**	-10.42 **	21.5	-1.86 **	29.95	-1.21 **	5.8*	-0.31 **
CG91 X CG45E	102.5	3.17 **	4.2	0.07 ns	271.4	9.50 **	24	1.36 *	32.15**	1.33 **	5.75	0.09 ns
CG91 X CG305	93.5	5.31 **	4.05	0.03 ns	243.15	0.92 ns	22.5	0.50 ns	31.15**	-0.12 ns	6.45**	0.22 **
CG150 X CG45SB	118.5**	7.02 **	4.35**	-0.06 ns	284.25**	3.78 **	24.5	-0.52 ns	32.2**	-0.78 **	6.75**	-0.19 *
CG150 X CG45E	71.5	-7.33 **	4.35**	0.05 ns	225.65	-7.85 **	23	-1.31 *	32.95**	0.31 **	6.9**	0.41 **
CG150 X CG305	68	0.31 ns	4.2	0.01 ns	217.9	4.07 **	25.5**	1.83 **	33.55**	0.46 **	6.85**	-0.21 **
CG62 X CG45SB	154**	6.02 **	3.95	-0.28 **	323.5**	7.85 **	23	-0.69 ns	28.65	0.55 **	5.35	-0.11 ns
CG62 X CG45E	108	-7.33 **	4.35**	0.23 **	251.9	-16.78 **	23.5	0.52 ns	26.65	-1.10 **	4.75	-0.26 **
CG62 X CG305	105.5	1.31 ns	4.05	0.05 ns	257.95	8.93 **	22.5	0.17 ns	28.75	0.55 **	5.95**	0.37 **
CG67 X CG45SB	161.5**	6.86 **	4.25*	0.15 *	342**	4.31 **	24.5	0.81 ns	28.2	0.19 *	5.3	0.17 *
CG67 X CG45E	120.5**	-1.50 ns	4.05	0.07 ns	284.5**	-6.21 **	25*	2.02 **	27.75	0.08 ns	4.65	-0.03 ns
CG67 X CG305	105.5	-5.36 **	3.65	-0.22 **	272.95	1.90 ns	19.5	-2.83 **	27.85	-0.27 **	5.1	-0.15 *
CG92 X CG45SB	112.5*	-4.48 **	3.95	-0.11 ns	275.45*	-3.60 **	22.5	-1.02 ns	28.55	-1.48 **	5.25	0.16 *
CG92 X CG45E	86.5	2.17 *	4.05	0.10 ns	244.05	11.97 **	23.5	0.69 ns	30.15	0.46 **	4.35	-0.29 **
CG92 X CG305	75.5	2.31 *	3.85	0.01 ns	204.05	-8.37 **	22.5	0.33 ns	31.15**	1.01 **	5.35	0.14 ns
CG163 X CG45SB	109	-13.48 **	4.15	-0.05 ns	282.4**	-14.89 **	23.5	-0.02 ns	31.35**	1.52 **	5.9**	0.34 **
CG163 X CG45E	87	-2.83 *	3.95	-0.13 ns	244.6	-5.71 **	21.5	-1.31 *	29.7	0.21 *	5.3	0.19 *
CG163 X CG305	95	16.31 **	4.15	0.18 *	251.25	20.60 **	23.5	1.33 *	28.2	-1.74 **	5.15	-0.53 **
Mean of hybrids	109.6		4.06		270.62		22.88		29.94		5.58	
CD 5 %	2.97		0.19		3.58		1.77		0.22		0.2	
CD 1 %	4.05		0.26		4.88		2.42		0.3		0.28	

* Significant at 5 % probability level, ** Significant at 1 % probability level.

Contd.....

Table 4. *Per se* performance and estimates of specific combining ability (*sca*) effects of hybrids for twelve characters in cotton

CROSS	Seed Index (g)		2.5 Span Length (mm)		Unifomity Ratio (%)		Fibre Strength (g/tex)		Micronaire		Elongation	
	MEAN	<i>Sca</i>	MEAN	<i>sca</i>	MEAN	<i>sca</i>	MEAN	<i>sca</i>	MEAN	<i>sca</i>	MEAN	<i>sca</i>
CG64 X CG45SB	12.75	-0.61 **	39.45**	0.62 **	47.15**	0.84 **	33.15**	0.42 **	3	-0.16 *	6.05**	0.66 **
CG64 X CG45E	12.55	-0.01 ns	37.4	-0.34 *	45.2**	-0.40 **	31.05**	-0.52 **	3.45*	0.30 **	5.15	-0.09 ns
CG64 X CG305	13.55**	0.62 **	36.55	-0.29 *	44.15	-0.44 **	29.75	0.10 ns	3.35	-0.14 *	5.15	-0.56 **
CG91 X CG45SB	13.75**	-0.11 ns	36.4	-0.96 **	44.45	-0.45 **	27.65	-0.55 **	3.65**	0.41 **	5.3	-0.34 **
CG91 X CG45E	12.35	-0.71 **	37.3	1.03 **	44.35	0.17 *	26.35	-0.69 **	3.15	-0.09 ns	5.75**	0.26 **
CG91 X CG305	14.25**	0.82 **	35.3	-0.07 ns	43.45	0.28 **	26.35	1.23 **	3.25	-0.32 **	6.05**	0.09 ns
CG150 X CG45SB	13.65**	0.04 ns	38.45**	0.94 **	43.15	-0.78 **	27.35	-0.88 **	3.45*	0.11 ns	5.75**	-0.18 *
CG150 X CG45E	13.5**	0.69 **	35.75	-0.67 **	42.6	-0.61 **	26.25	-0.82 **	3.05	-0.29 **	6.15**	0.37 **
CG150 X CG305	12.45	-0.73 **	35.25	-0.27 *	43.6	1.39 **	26.85	1.70 **	3.85**	0.18 *	6.05**	-0.20 **
CG62 X CG45SB	13.25**	-0.31 **	39.8**	0.10 ns	45.25**	0.37 **	35.7**	1.72 **	2.8	-0.08 ns	5.25	0.27 **
CG62 X CG45E	13.05*	0.29 **	38.95**	0.35 *	44.75**	0.59 **	33.35**	0.53 **	2.75	-0.12 ns	4.45	-0.38 **
CG62 X CG305	13.15**	0.02 ns	37.25	-0.45 **	42.2	-0.96 **	28.65	-2.25 **	3.4*	0.20 **	5.4	0.10 ns
CG67 X CG45SB	13.15**	0.06 ns	39.3**	-0.25 ns	45.75**	0.19 **	34.55**	0.24 *	2.95	0.16 *	4.25	-0.43 **
CG67 X CG45E	12.05	-0.24 **	38.9**	0.45 **	44.25	-0.60 **	34**	0.85 **	2.6	-0.19 *	4.75	0.22 **
CG67 X CG305	12.85	0.19 *	37.35	-0.20 ns	44.25	0.41 **	30.15	-1.08 **	3.15	0.03 ns	5.2	0.20 **

Contd.....

CG92 X CG45SB	13.15**	1.01 **	38.95**	0.10 ns	44.25	-0.88 **	32.2**	-0.98 **	2.65	-0.39 **	5.25	-0.18 *
CG92 X CG45E	11.1	-0.24 **	37.45	-0.30 *	45.25**	0.84 **	32.85**	0.83 **	3.25	0.21 **	4.85	-0.43 **
CG92 X CG305	10.95	-0.76 **	37.05	0.20 ns	43.45	0.04 ns	30.25	0.15 ns	3.55**	0.18 *	6.35**	0.60 **
CG163 X CG45SB	12.95	-0.08 ns	38.3**	-0.56 **	46.65**	0.70 **	31.15**	0.02 ns	3.25	-0.04 ns	5.45	0.19 **
CG163 X CG45E	12.45	0.22 **	37.25	-0.52 **	45.25**	0.02 ns	29.8	-0.17 ns	3.45*	0.16 *	5.15	0.04 ns
CG163 X CG305	12.45	-0.15 ns	37.95	1.08 **	43.5	-0.72 **	28.2	0.15 ns	3.5**	-0.12 ns	5.35	-0.23 **
Mean of hybrids	12.83		37.64		44.42		30.27		3.21		5.39	
CD 5 %	0.21		0.37		0.18		0.28		0.19		0.18	
CD 1 %	0.29		0.5		0.24		0.38		0.26		0.25	

*Significant at 5 % probability level, ** Significant at 1 % probability level.

Table 5. Relationship between *per se*, *sca* effects and Standard heterosis for twelve characters in cotton

Sl.No	Character	Hybrids	<i>per se</i>	Hybrids	<i>sca</i> effects	Hybrids	Standard Heterosis (%)	Hybrids selected as whole
1.	Number of bolls per plant	CG64 X CG45SB	166.50	CG163 X CG305	16.31	CG64 X CG45SB	129.66	CG64(24.90)X CG45SB (25.48)
		CG67 X CG45SB	161.50	CG64 X CG45E	13.67	CG67 X CG45SB	122.76	
		CG62 X CG45SB	154.00	CG150 X CG45SB	7.02	CG62 X CG45SB	112.41	CG67 (19.57) X CG45SB (25.48)
2.	Boll Weight (g)	CG64 X CG45SB	4.40	CG64 X CG45SB	0.45	CG64 X CG45SB	3.53	CG64 (-0.22) X CG45SB (0.11)
		CG150 X CG45SB	4.35	CG62 X CG45E	0.23	CG150 X CG45SB	2.35	
		CG150 X CG45E	4.35	CG163 X CG305	0.18	CG150 X CG45E	2.35	CG150(0.24) X CG45SB (0.11)
3.	Seed cotton yield per plant (g)	CG64 X CG45SB	353.40	CG163 X CG305	20.60	CG64 X CG45SB	49.81	CG64 (31.94) X CG45SB (37.87)
		CG67 X CG45SB	342.00	CG64 X CG45E	15.09	CG67 X CG45SB	44.98	
		CG62 X CG45SB	323.50	CG64 X CG45SB	12.96	CG62 X CG45SB	37.13	CG67(29.19) X CG45SB (37.87)
4.	Number of seeds per boll	CG64 X CG45SB	25.50	CG64 X CG45SB	3.31	CG64 X CG45SB	2.00	CG64 (-1.38) X CG45SB (0.69)
		CG150 X CG305	25.50	CG67 X CG45E	2.02	CG150 X CG305	2.00	CG150 (1.45)X CG305(-0.67)
		CG67 X CG45E	25.00	CG150 X CG305	1.83			
5.	Ginning outturn (per cent)	CG150 X CG305	33.55	CG163 X CG45SB	1.52	CG150 X CG305	7.02	CG150 (2.96) X CG305 (0.19)
		CG150 X CG45E	32.95	CG91 X CG45E	1.33	CG150 X CG45E	5.10	
		CG150 X CG45SB	32.20	CG64 X CG45SB	1.20	CG150 X CG45SB	2.71	CG150 (2.96) X CG45E (-0.26)
6.	Lint Index (g)	CG64 X CG45SB	31.20					
		CG150 X CG45E	6.90	CG150 X CG45E	0.41	CG150 X CG45E	22.12	CG150 (1.26) X CG45E (-0.34)
		CG150 X CG305	6.85	CG62 X CG305	0.37	CG150 X CG305	21.24	
7.	Seed Index (g)	CG150 X CG45SB	6.75	CG163 X CG45SB	0.34	CG150 X CG45SB	19.47	CG150 (1.26) X CG305(0.23)
		CG64 X CG45SB	5.45					
		CG91 X CG305	14.25	CG92 X CG45SB	1.01	CG91 X CG305	13.10	CG91 (0.62) X CG305 (-0.02)
8.	2.5 Span Length (mm)	CG91 X CG45SB	13.75	CG91 X CG305	0.82	CG91 X CG45SB	9.13	
		CG150 X CG45SB	13.65	CG150 X CG45E	0.69	CG150 X CG45SB	8.33	CG91(0.62) X CG45SB (0.41)
		CG62 X CG45SB	39.80	CG163 X CG305	1.08	CG62 X CG45SB	8.45	CG 62 (1.03) X CG45SB (1.03)
9.	Uniformity Ratio (per cent)	CG64 X CG45SB	39.45	CG91 X CG45E	1.03	CG64 X CG45SB	7.49	
		CG67 X CG45SB	39.30	CG150 X CG45SB	0.94	CG67 X CG45SB	7.08	CG 64(0.16) X CG45SB (1.03)
		CG64 X CG45SB	47.15	CG150 X CG305	1.39	CG64 X CG45SB	9.65	CG64 (1.08) X CG45SB (0.81)
10.	Fibre Strength (g/tex)	CG163 X CG45SB	46.65	CG64 X CG45SB	0.84	CG163 X CG45SB	8.49	
		CG67 X CG45SB	45.75	CG92 X CG45E	0.84	CG67 X CG45SB	6.40	CG163 (0.71) X CG45SB (0.81)
		CG62 X CG45SB	35.70	CG62 X CG45SB	1.72	CG62 X CG45SB	30.53	CG62 (2.30) X CG45SB (1.41)
11.	Micronaire	CG67 X CG45SB	34.55	CG150 X CG305	1.70	CG67 X CG45SB	26.33	
		CG67 X CG45E	34.00	CG91 X CG305	1.23	CG67 X CG45E	24.31	CG67(2.63)X CG45SB (1.41)
		CG64 X CG45E	33.35			CG64 X CG45E	21.94	
12.	Elongation	CG64 X CG45SB	33.15			CG64 X CG45SB	21.21	
		CG150 X CG305	3.85	CG91 X CG45SB	0.41	CG150 X CG305	18.46	CG150 (0.24)X CG305(0.22)
		CG91 X CG45SB	3.65	CG64 X CG45E	0.30	CG91 X CG45SB	12.31	
12.	Elongation	CG92 X CG305	3.55	CG92 X CG45E	0.21	CG92 X CG305	9.23	CG91(0.14)X CG45SB (-0.11)
		CG92 X CG305	6.35	CG64 X CG45SB	0.66	CG92 X CG305	12.39	CG92(0.10)X CG305(0.26)
		CG150 X CG45E	6.15	CG92 X CG305	0.60	CG150 X CG45E	8.85	
		CG64 X CG45SB	6.05	CG150 X CG45E	0.37	CG64 X CG45SB	7.08	CG150(0.60)X CG45E(-0.21)

*Figures in parenthesis indicates *sca* effects

Marked negative *sca* effect for 2.5 per cent span length in the cross CG 67 x CG 45 SB could be attributed to the lack of co-adaptation between favorable alleles of the parents involved. Gururaj Rao *et al.* (1977) reported that this was probably due to mutual cancellation of components of heterosis. The results are in agreement with the research findings of Patel *et al.* (2012) and Sekhar *et al.* (2012).

In the present investigation, the parent CG 64, CG 67 and CG 45 SB were found to be promising due to its high yield potential as well as significant and positive general combining ability effects for yield and its attributes and for 2.5 per cent span length and fibre strength. Therefore, these three parents could be effectively used for hybrid breeding programme as well as introgression breeding. The cross combinations having high mean yield, high heterosis and desirable *sca* effects is of immense importance for hybrid cotton breeding programme. For seed cotton yield and major yield attributes as well as important fibre quality parameters the CG 64 x CG 45SB and CG 67 x CG 45 SB showed positive and significant, *per se*, standard heterosis and *sca* effects. Consequently, these two hybrids appeared promising for commercial release after thorough testing. The results are in conformity with the reports of Maisurua *et al.* (2007) and Sekhar *et al.* (2012).

REFERENCES

- AICCIP annual Report. 2013-14. All India Coordinated Cotton Improvement Project. Coimbatore, Tamilnadu, India.
- Amir, S., Farooq, J., Ameer, B., Khan, S.H and Saleem, M.F. 2012. Genetic studies of earliness in *Gossypiumhirsutum* L. *International Journal of Agriculture, Veterinary and Medical Sciences*, 6 (3): 189-207
- Gururaj Rao, M. R., Hiremath, K. G. and Virupakshappa, K. 1977. Genetic analysis of ginning and fibre properties in upland cotton *G. hirsutum* L. II. Combining ability studies in respect of six quantitative characters. *Mysore J. Agric. Sci.*, 2(4): 453-457.
- Kempthorne, O. 1957. An introduction to genetic statistics. John Wiley and Sons Inc., New York. 458-471.
- Liang, G.H., Reddy, C.R. and Dayton, A.D. 1971. Heterosis, inbreeding depression and heritability estimates in a systematic series of grain sorghum genotypes. *Crop Science*, 12: 409-411.
- Maisuria, A.T., Patel, J.C., Patel, K.G., Patel, D.H and Chhimpi, B.G. 2007. Heterosis for yield and its contributing characters in gms based Asiatic cottons. *Journal of Indian Society for Cotton Improvement*, 32 (1): 30-40.
- Patel, N.A., Patel, B.N., Bhatt, J.P and Patel, J.A. 2012. Heterosis and combining ability for seed cotton yield and component traits in inter-specific cotton hybrids (*Gossypiumhirsutum* L. x *Gossypiumbarbadense* L.). *Madras Agricultural Journal*, 99 (10-12): 649-656.
- Pathak, R. S. and Kumar, P. 1975. Combining ability studies in upland cotton *G. hirsutum*L. *Zeitschrift fur Pflanzenchtung*, 75(4): 297-310.
- Rao, P.J.M. and Gopinath, M. 2013. Variability and association studies for yield and yield components in upland cotton (*Gossypiumhirsutum* L.) under chalka soils.
- Sekhar, L., Khadi, B.M., Rajesh, S.P., Katageri, I.S., Vamadevaiah, H.M., Chetti, M.B and Nadaf, H.L. 2012. Study of heterosis in thermo sensitive genetic male sterility (TGMS) based diploid cotton hybrids for yield, yield component and fibre quality characters. *Karnataka Journal of Agricultural Sciences*, 25 (3): 313-321.
- Shull, G.H. 1948. What is heterosis? *Genetics*, 33: 439-446
- Simon, S.Y., Kadams, A.M. and Aliyu, B. 2013. Combining ability analysis in F1 hybrids of cotton (*Gossypium species* L.) by diallel method in North eastern Nigeria. *Greener Journal of Agricultural Sciences*, 3(2): 090-096.
- Sprague, G. F. and Tatum, L. M. 1942. General versus specific combining ability in single crosses of corn. *Agron. J.*, 34: 923-932.
- Vineela, N., Murthy, J.S.V.S., Kumar, P.V.R and Kumari, S.R. 2012. Combining ability estimates for yield and fibre quality traits in line x tester crosses of intra *hirsutum* hybrids of cotton (*Gossypiumhirsutum* L.). *The Andhra Agricultural Journal*, 59 (4): 545-555.
