



ISSN: 0975-833X

Available online at <http://www.journalcra.com>

International Journal of Current Research
Vol. 11, Issue, 05, pp.3641-3645, May, 2019

DOI: <https://doi.org/10.24941/ijcr.34801.05.2019>

INTERNATIONAL JOURNAL
OF CURRENT RESEARCH

RESEARCH ARTICLE

A CROSS SECTIONAL OBSERVATIONAL STUDY ON NON-COMMUNICABLE DISEASES AMONG ELDERLY POPULATION IN BLOCK HAZRATBAL, DISTRICT SRINAGAR

^{1,*}Dr. Umar Nazir, ¹Dr. Rouf Hussain Rather, ²Dr. Shazia Benazir and
³Prof. Mohammad Salim Khan, S.

¹Demonstrator, Department of Community Medicine, Government Medical College, Karanagar Srinagar

²Senior Resident at Department of Microbiology SKIMS Soura, Srinagar

³Professor and Head Department of Community Medicine, Govt. Medical College, Srinagar

ARTICLE INFO

Article History:

Received 20th February, 2019
Received in revised form
14th March, 2019
Accepted 10th April, 2019
Published online 30th May, 2019

Key Words:

Non-communicable
Diseases (NCDs), Elderly.

ABSTRACT

Majority of Elderly population aged 60 years and above will be in the developing countries as their population will reach 1.2 billion by the year 2025(1). Major health issues which they face in this age group are Non-communicable diseases like hypertension, coronary artery disease, strokes, diabetes mellitus, obesity, cancers, cataracts, osteoarthritis, osteoporosis, chronic obstructive airway disease, benign hyperplasia of prostate, Alzheimer's disease, Parkinson's disease, Senile Dementia and depression (2). It has been estimated by W.H.O that by 2020, heart disease and stroke will become the leading causes of death and disability worldwide, and the number of fatalities projected to increase to more than 24 million by 2030 (3). Burden of diseases is shifting from communicable to non-communicable in the developing countries of Asia because of the Demographic and epidemiological transitions taking place in this part of the world India's poor are at heightened risk of acquiring NCDs owing to high rate of smoking, tobacco use and high salt intake. So this paper analyses the links between sociodemographic correlates and non-communicable diseases in block Hazratbal, district Srinagar. **Objectives:** 1) To estimate the prevalence of NCDs among study population. 2) To find out the socio-demographic correlates of NCDs in the study population. **Study design:** A Community Based, Cross sectional study. **Study population:** Elderly (≥ 60 years) persons residing in selected geographical area. **Results:** In this study, majority of the elderly study population were suffering from Hypertension/Ischemic/Other heart diseases (52.2%) followed by COPD (11.4%), Arthritis/other Musculoskeletal problems (10.6%), Diabetes mellitus (10.4%). Gastritis (9.8%), Hypothyroidism (9.6%), Benign prostrate hypertrophy (9.4%). The main reason for high prevalence of Hypertension/Ischemic heart disease could be that we take lots of salty products in our daily diet. Other causes could be living in a conflict area or genetic susceptibility. Most of elderly study subjects 446(89.9%) were on drugs for their chronic diseases.

*Corresponding author: Dr. Umar Nazir

Copyright ©2019, Umar Nazir et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Dr. Umar Nazir, Dr. Rouf Hussain Rather, Dr. Shazia Benazir, Prof. Mohammad Salim Khan, S. 2019. "A cross sectional observational study on Non-Communicable Diseases among Elderly population in Block Hazratbal, District Srinagar.", *International Journal of Current Research*, 11, (05), 3641-3645.

INTRODUCTION

Majority of Elderly population aged 60 years and above will be in the developing countries as their population will reach 1.2 billion by the year 2025 (Arokiasamy, 1997). Major health issues which they face in this age group are Non-communicable diseases like hypertension, coronary artery disease, strokes, diabetes mellitus, obesity, cancers, cataracts, osteoarthritis, osteoporosis, chronic obstructive airway disease, benign hyperplasia of prostate, Alzheimer's disease, Parkinson's disease, Senile Dementia and depression (Burke, 2000). It has been estimated by W.H.O that by 2020, heart disease and stroke will become the leading causes of death and disability worldwide, and the number of fatalities projected to increase to more than 24 million by 2030 (Joy Kumar Chakma, 2014).

Burden of diseases is shifting from communicable to non-communicable in the developing countries of Asia because of the Demographic and epidemiological transitions taking place in this part of the world(6). In India alone, rapid changes in the country's Socio-cultural and lifestyles have caused NCD's to become responsible for two-thirds of the total morbidity burden and about 53% of total deaths (Joy Kumar Chakma et al., 2014). The simultaneous occurrence of more than one health condition in the same individual especially in the older adults is increasingly becoming the norm.

Rationale of the study

India's Geriatric population are at heightened risk of acquiring NCDs owing to high rate of smoking, tobacco use, physical inactivity, unhealthy diet, urbanization, stress and high salt

intake especially in this part of the world.(5)So this study throws some light to know about the Magnitude of NCD's among geriatric population in Block Hazratbal which is the field practice area of Government Medical College, Srinagar.

OBJECTIVES

- To estimate the prevalence of NCD's among geriatric population in Block Hazratbal, District Srinagar.
- To find out the Socio-demographic correlates of NCD's in the study population.

MATERIALS AND METHODS

- Study design:** A Cross-sectional, Community Based study.
- Study population:** (≥ 60 years of both genders)
- Inclusion Criteria:**

Permanent Residents of Kashmir Valley

- Age 60 years and above.

Exclusion Criteria

Geriatric population who refuse to give informed consent for the study.

- Study area:** Block Hazratbal of District Srinagar
- Study duration:** From 1st April 2015 – 30th September 2016

Sample size calculation

$N = (1.96)^2 pq / L^2$
496 elderly were selected.

Sample Plan: As shown in flow-chart below

Sample Method: Multi-stage Random Sampling technique was used.

In the first stage 50% health centres were randomly chosen from each zone using **lottery method**

In the second stage, requisite number of study units were selected from each selected health center **randomly**.

Ethical Consideration: All the selected subjects were informed about the objectives of the study. A proper written informed consent was taken from the selected subjects who agreed to participate in the study.

Procedure: The identified 496 elderly subjects were interviewed in their local language and they were examined by using a pre-tested, semi-structured, study questionnaire. The questionnaire was divided into two parts. The first part comprised of the socio-demographic information. In second part the diagnosis was obtained from written Medical records

of Registered Medical Practitioners (Both Public & Private Sector) available with the study subjects.

OBSEVATIONS AND RESULTS

Table 1. Distribution of study population as per their gender

Gender	No. of elderly (n)	Percentage (%)
Female	269	54.2
Male	227	45.8
Total	496	100.0

Table 2. Distribution of study population as per their marital status

Marital Status	No. of elderly (n)	Percentage (%)
Currently Married	401	80.8
Widow/Widower	92	18.5
Separated	1	0.2
Divorced	1	0.2
Never Married	1	0.2
Total	496	100.0

Table 3. Distribution of study population as per their educational qualification

Education	No. of elderly (n)	Percentage (%)
Illiterate	358	72.2
Primary	47	9.5
Middle	41	8.3
Secondary	24	4.8
Higher Secondary	7	1.4
Graduate/Post graduate	18	3.6
Professional	1	0.2
Total	496	100.0

Table 4. Distribution of study subjects according to socio-economic status

Socio-Economic status (Kuppuswami scale 2016)	No. of elderly (n)	Percentage (%)
Cat-I (Upper)	5	1.0
Cat-II (Upper Middle)	179	36.1
Cat-III (Lower Middle)	107	21.6
Cat-IV (Upper Lower)	205	41.3
Cat-V (Lower)	0	0.0
Total	496	100.0


Table 5. Distribution of study subjects according to direct effect of conflict on their lives in the region

Life in conflict area	No. of elderly (n)	Percentage (%)
Not effected	438	88.3
Effected	58	11.7
Total	496	100.0

Table 6. Distribution of Non Communicable Diseases in Study Subjects

Morbidity	No. of elderly(n=496)	Percentage (%)
Hypertension / IHD	259	52.2
DEPRESSION	156	31.5
COPD	57	11.4
Musculoskeletal problems	53	10.6
Diabetes mellitus	52	10.4
Gastritis	49	9.8
Hypothyroidism	48	9.6
Benign prostrate hypertrophy	47	9.4
Osteoporosis	17	3.4
Cataracts	11	2.2
Sensory Neural Hearing Loss	10	2.0
Cancers	06	1.2

*Males only.



Prevalence of Non-communicable Diseases was found 52.2%.

Table 7. Distribution of NCD's according to Gender

NCD's	Gender		Total
	Male	Female	
Present	103(39.5%)	156(60.5%)	259 (100%)
Absent	124(52.4%)	113(47.6%)	237 (100%)
Total	227(45.8%)	269(54.2%)	496(100%)

Table 8. Distribution of NCD's according to different age groups

AGE (In Years)	NCD's		Total n (%)
	PRESENT n (%)	ABSENT n (%)	
60 - 69	108(48.7%)	114(51.35%)	222(100.0%)
70 - 79	126(53.6%)	109(46.4%)	235(100.0%)
80 & above*	25(64.1%)	14(35.9%)	39(100.0%)
Total	259(52.2%)	237(47.8%)	496(100%)

Table 9. Distribution of NCD's according to educational status in the study group

Educational status	NCD's		Total n (%)
	PRESENT n (%)	ABSENT n (%)	
Illiterate	229 (64%)	129 (36%)	358(100.0%)
Literate*	30 (21.7%)	108(78.3%)	138(100.0%)
Total	259(52.2%)	237 (47.8%)	496(100.0%)

Table 10. Distribution of NCD's according to marital status in the study group

Marital Status	NCD's		Total n (%)
	PRESENT n (%)	ABSENT n (%)	
Currently Married	170(47.2%)	190(52.77%)	360(100.0%)
Widow/widower & Single	89(65.4%)	47(34.55%)	136(100.0%)
Total	259(52.2%)	237(47.8%)	496(100.0%)

Table 11. Distribution of NCD's according to Socio economic category in study subjects

Socio-economic Category	NCD's		Total n (%)
	PRESENT n (%)	ABSENT n (%)	
Upper/Upper Middle*	40(30.3%)	92(69.69%)	132(100.0%)
Lower Middle	66(51.16%)	63(48.8%)	129(100.0%)
Upper Lower	153(65.10%)	82 (34.89%)	235(100.0%)
Total	259(52.2%)	237(47.8%)	496(100.0%)

X²=60.035. P<0.001**

Table 12. Distribution of NCD's according to Economic dependence

Economic dependency	NCD's		Total n (%)
	PRESENT n (%)	ABSENT n (%)	
INDEPENDENT	30(23.62%)	97 76.37%)	127(100.0%)
Partially dependent	84(70.58%)	35 (29.42%)	119(100.0%)
Totally dependent	145(58.00%)	105 (42.00%)	250 (100.0%)
Total	259 (52.2%)	237 (47.8%)	496 (100.0%)

X²=52.199. P<0.001

Table 13. Relationship of NCD's with dependency for the activities of daily living

Dependency for the Activities of Daily Living	NCD's		Total n (%)
	PRESENT n (%)	ABSENT n (%)	
Independent	180(45.56%)	215(54.4%)	395(100.0%)
Partially dependent/ totally dependent*	79 (78.2%)	22(21.7%)	101(100.0%)
Total	259(52.2%)	237(47.8%)	496(100.0%)

X²=21.997. P<0.001

Table 14. Relationship of NCD's with Depression in study subjects

Depression	NCD's		Total N (%)
	PRESENT n (%)	ABSENT n (%)	
Present	156(61.4%)	98(38.6%)	254 (100.0%)
Absent	103(42.6 %)	139(57.4%)	242(100.0%)
Total	259(52.2 %)	237(47.8%)	496(100.0%)

X²=9.962. P<0.002

Table 15. Relationship of NCD's with Tobacco consumption status in the elderly study subjects

Tobacco Consumption	NCD's		Total n (%)
	PRESENT n (%)	ABSENT n (%)	
Current Tobacco Consumer	183(69.58%)	80(30.41%)	263(100.0%)
Ex / Non Tobacco Consumer	76(32.61%)	157(67.38%)	233(100.0%)
Total	259(52.2%)	237(47.8%)	496(100.0%)

X²=2.753Chi-square test**P<0.005**

Table 16. Relationship of NCD's with drug intake of study subjects for their chronic diseases

DRUG INTAKE	NCD's		Total n (%)
	Present n (%)	Absent n (%)	
Present	210(55.7%)	167(44.3%)	377(100%)
Absent	49 (41.1%)	70(58.8%)	119(100%)
Total	259(52.2%)	237(47.8%)	496 (100%)

X²= 11.869. P <0.001

The prevalence of NCD was highest (55.7%) among the study subjects who were on drugs for their chronic diseases. The difference was found statistically significant (P<0.001).

DISCUSSION

More than 52.2% of the older adults in our study had at least one NCD which is a huge challenge for the healthcare system in India with limited capacity for managing NCDs. In our study it was seen that Hypertension/IHD had a prevalence of 52%, depression 31.5%, COPD 11.4%, Musculoskeletal problems 10.6%, followed by Diabetes mellitus 10.4%. A study conducted by Joy Kumar Chakma & Sanjay Gupta on "Lifestyle and Non-Communicable Diseases: A double edged sword for future India" showed that in India, 53% of the deaths in 2008 were due to NCDs (WHO). The cardiovascular diseases (CVDs) alone account for 24 percent of all deaths. The anticipated cumulative loss of national income due to NCDs mortality for India for 2006-2015 will be USD237 billion. By 2030, this productivity loss is expected to double (Joy Kumar Chakma, 2014). In our study NCDs were more common in females 156(60.5%) as compared to males where it

has a prevalence of 103(39.5%).The difference was also found statistically significant.($p<0.005$).One of the studies in Chandigarh found out that female elderly were more prone to morbidity (Swami et al., 2002) (National Program for Prevention and Control of Diabetes, 2008-2009). In our study NCDs were highest in age group of 80 years and above 25(64.1%) and lowest in the age group of 60-69years where it was 108 (48.7%).The difference was also found statistically significant. ($p<0.005$)The results of similar studies are also consistent with our study where the incidence of Multi-morbidities positively associated with age, with an increase in age the proportion of no morbidity to one morbidity and multi morbidity increased many folds . (Arokiasamy, 1997; Chirinda, 2015). In our study NCDs were more common in study subjects who were illiterate 221(64%) as compared to literates where it was found only 30(21.7%).The difference was also found statistically significant ($p<0.005$).

Our study found that the geriatric population who were widow/widower and single had NCDs 89(65.4%) more common than who were currently married 170(47.2%).The difference was also found statistically significant ($p<0.005$). Our study found that the geriatric population who were belonging to upper lower class had NCDs 153(65.1%) more common than who were belonging to upper/upper middle class 40 (30.3%).The difference was also found statistically significant. $X^2=60.035(p<0.005)$ In another study the association between socio-economic status and prevalence of specific chronic diseases is well established (Marmot, 2005)A study on multi-morbidity in Australia found that 85% of 70+ year elderly have multi-morbidity and the prevalence is higher among elderly, with obese, female, in low socio-economic status, living alone and less educate, leading to negative impact on quality of life (Walker, 2007) (Marmot, 2005; Walker, 2007). In our study NCDs was highest in the geriatric population who were partially dependent economically on others 84(70.5%) and lowest who were economically independent 30 (23.6%). The difference was also found statistically significant. $x^2=52.19(p<0.001)$.

In another study where age, state of economic independence and lifestyle indicators are more important predictors of multi-morbidity among study population was studied (Banjare & Pradhan, 2014). In our study NCDs was highest in the geriatric population who were partially/ totally dependent for the activities of daily living 79 (78.2%) and lowest who were independent for the activities of daily living 180 (45.5%). The difference was also found statistically significant. $x^2=21.997(p<0.001)$. In our study NCDs were more common in study subjects who had depression 156 (61.4%) as compared to study subjects where depression was found absent 103(42.6%). The difference was also found statistically significant. $X^2=9.962$ ($p<0.002$). In our study NCDs was highest in the geriatric population who were tobacco consumers 183(69.5%) and lowest who were Ex- Nonsmokers 76 (32.6%).The difference was also found statistically significant. $x^2=2.753(p<0.005)$. In our study NCDs was more common in the geriatric population who were on drugs for their chronic morbidities 210(55.7%) as compared who were not on drugs for their chronic ailments 49 (41.1%).The difference was also found statistically significant. $x^2=11.869(p<0.001)$.

Conclusion

- Overall prevalence of NCD's was higher among the Geriatric population in this part of the India.
- NCD's was found more in females as compared to males.
- Major NCD's which were found are Hypertension/CVD, depression, COPD, Diabetes Mellitus and degenerative disorders.
- Strong predictors on the prevalence of NCD's in the elderly were .advanced age, illiteracy, marital status, socio-economic class, family pattern, economic dependence, dependence for activities of daily living, Tobacco consumption, drug intake for chronic ailments.

Recommendation

- Mass Screening for NCD's in Geriatric population should be done regularly in primary care health delivery system.
- A comprehensive strategy for the prevention and control of NCDs must integrate public health actions to minimize risk factor exposure at the level of the population and reduce risk at the level of individuals at high risk
- Need of the hour is to strengthening the existing "package" of services for elderly under various initiatives and programs by means of IEC (Information Education & Communication) and BCC (Behavioral Change Communication) especially to bring a social change targeting towards the Geriatric population.
- Health services need to be strengthened accordingly with increased emphasis on key service utilization determinants like service availability and accessibility.
- The findings of our study could guide Health Policy Makers to devise and implement effective and timely NCD based health interventions for the Geriatric population, in order to decrease geriatric NCD's and develop a comprehensive strategy for its early diagnosis and early management without getting in complications.

REFERENCES

- Arokiasamy JT. 1997. Malaysia's Ageing Issues (Editorial). *Med J Malaysia* 52:197-201.
- Banjare, P., & Pradhan, J. 2014. Socio-economic inequalities in the prevalence of multi- morbidity among the rural elderly in Bargarh district of Odisha (India). *PloS one*, 9(6), e97832.
- Burke MM, Laramie JA. 2000. Primary care of the older adult. 1stEdition. USA: Mosby Inc.
- Chirinda, W., et al. 2015. "Self-reported prevalence of chronic non-communicable diseases and associated factors among older adults in South Africa."
- Economics of Non-Communicable Diseases in India A report by the World Economic Forum and the Harvard School of Public Health. {Online} 2014 November.
- Joy Kumar Chakma, Sanjay Gupta.Lifestyle and Non-Communicable Diseases: A double edged sword for future India. *Indian journal of community health* [Serial Online] 2014; 26:4. Available from:<http://www.iapsmupuk.org/journal/index.php/IJCH/article/view/716>

- Marmot, M. 2005. Social determinants of health inequalities. *The Lancet*, 365(9464), 1099- 1104.
- National Program me for Prevention and Control of Diabetes, Cardiovascular Disease and Stroke. Manual for medical officers. 2008-2009.
- Swami, H. M., Bhatia, V., Dutt, R., & Bhatia, S. P. S. 2002. A community based study of the morbidity profile among the elderly in Chandigarh, India. *Bahrain Medical Bulletin*, 24(1), 16-20.
- Walker, A. E. 2007. Multiple chronic diseases and quality of life: patterns emerging from large national sample, Australia. *Chronic Illness*, 3(3), 202-218.
- World Health Organization 2011. Cause-specific mortality, 2008: WHO region by country. Geneva: WHO.
