



ISSN: 0975-833X

RESEARCH ARTICLE

PRIMARY PRODUCTIVITY OF BAY OF BENGAL AT PARADIP IN ODISHA, INDIA

*Rajalaxmi Mohanty and Patra, A. K.

Fishery Science Research Laboratories, P.G. Department of Zoology, Utkal University, Vanivihar,
Bhubaneswar, Odisha, India

ARTICLE INFO

Article History:

Received 15th September, 2014
Received in revised form
20th October, 2014
Accepted 19th November, 2014
Published online 30th December, 2014

Key words:

Primary Productivity,
Bay of Bengal,
Paradip,
Odisha.

ABSTRACT

The primary productivity of Bay of Bengal at Paradip has been assessed both spatially and seasonally. The maximum GPP was noted during summer season and minimum in rainy season. The annual mean GPP varied from $0.077 + 0.008 \text{ g}^c \text{ m}^3 \text{ h}^{-1}$ to $0.996 + 0.16 \text{ g}^c \text{ m}^3 \text{ h}^{-1}$. The NPP value varied from $0.016 + 0.002 \text{ g}^c \text{ m}^3 \text{ h}^{-1}$ to $0.845 + 0.072 \text{ g}^c \text{ m}^3 \text{ h}^{-1}$ and showed an increasing trend from the month of September 2011 to February, 2012. The community respiration (CR) ranged from $0.027 + 0.003 \text{ g}^c \text{ m}^3 \text{ h}^{-1}$ to $0.499 + 0.046 \text{ g}^c \text{ m}^3 \text{ h}^{-1}$ and exhibited a systematic seasonal variation with maximum and minimum value during summer and winter respectively. The Correlation coefficients (R) of NPP, GPP and CR with certain physicochemical parameters have also been established.

Copyright © 2014 Rajalaxmi Mohanty and Patra, A. K. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

The primary productivity of a water body is the manipulation of its biological production. It forms the basis of the ecosystem functioning, (Odum, 1971). It plays an important role in energy and organic matters available to the entire biological community (Ahmed *et al.*, 2005). Primary productivity at each level can be distinguished further into gross primary production i. e the total amount of organic matters produced and net primary production or the amount of organic matter produced of a particular level (Yeragi and Shaikh, 2003). The estimation of primary productivity is predicted on the relationship between oxygen evolution and carbon fixation (Dash *et al.*, 2011). Primary productivity varies from fresh-water to estuarine and from estuarine to marine water body (Dash *et al.*, 2011). Most of the organic matters of an aquatic ecosystem are produced within the water by phyto-plankton (= primary producer) which is utilized by the consumer (Goldman and Wetzel, 1963). The area under study is Bay of Bengal at Paradip in Odisha. The Paradip beach stretches nearly 35 Km. along the Bay of Bengal and it is a noted visiting place of national and international repute. Further, it is an important fish landing centre of varieties of edible and commercially as well as economically important fin-fish, shall fish species. The drastic reduction is marine edible fin-fish and shall fish fauna and diversity along the Paradip sea-beach calls for immediate analysis and intermation.

*Corresponding author: Rajalaxmi Mohanty

Fishery Science Research Laboratories, P.G. Department of Zoology,
Utkal University, Vanivihar, Bhubaneswar, Odisha, India

MATERIALS AND METHODS

For the analysis of primary productivity Light and Dark bottle method is recommended by 'Vollenweiden'(1969), was followed. Gaarder and Gran (1927) first proposed the technique of using light and dark bottles and winkler's titration to measure the production and consumption of oxygen. Water samples were collected in triplicate in the middle of each month between 10:00 AM to 12:00 Noon. The sample in the first bottle was used immediately to determine the initial level of dissolved oxygen content following modified winkler's volumetric method (APHA, 1998). Dissolved Oxygen, values obtained in the present study were converted to carbon values by multiplying with the factor 0.375 (Odum, 1956). The second bottle was pointed with black color (dark bottle) to prevent light and hence serve as control to measure respiration. The third bottle (light) was treated as a test to measure the net production. The last two bottles were incubated under water in the emphatic zone for a period of six hours by suspending it in water after which, dissolved oxygen content of each bottle was estimated.

Gross primary production (GPP) and the net primary production (NPP) were calculated in the following manner.

$$\text{GPP (O}_2 \text{ mg l}^{-1} \text{ + h}^{-1} \text{) = (DI-Di)/h}$$

$$\text{NPP (O}_2 \text{ mg l}^{-1} \text{ + h}^{-1} \text{) = (DI-Di)/h}$$

Community Respiration

Community Respiration $O_2 \text{ mg } C^{-1} + h^{-1} = (D1-Di)/h$

Where, D1 = $O_2 \text{ (mg } C^{-1}h^{-1})$ in the higher bottle

Di = $O_2 \text{ (mg } C^{-1}h^{-1})$ in the initial bottle

Dd = $O_2 \text{ (mg } C^{-1}h^{-1})$ in the dark bottle

h = Duration of exposure in hours.

The hourly rate can be converted to daily rates by multiplying with duration of sunshine on that day. Oxygen values ($\text{mg } C^{-1}$) were converted to carbon value by applying the equation suggested by Thomas *et al.*, (1980).

Primary production (gc) = $(O_2 \text{ (mg } C^{-1}) \times 0.375) / PQ$
Where PQ = 1.25

PQ represents respiratory gradient = respiration / photosynthesis and a comprised value of 1-25 was used which represents metabolism of sugar, fat and proteins. The value 0.375 represents a constant to convert oxygen value to carbon value (Thomas *et al.*, 1980).

RESULTS AND DISCUSSION

The seasonal variation of gross primary productivity (GPP), net primary productivity (NPP) and community respiration (CR) of Bay of Bengal at Paradip is given in Table 1.

Table 1. Mean Seasonal Variations in GPP, NPP and Community Respiration (mean \pm SD) in $g^c m^3 h^{-1}$ at Paradip

Season	GPP	RES	NPP
Summer	0.800 \pm 0.091	0.243 \pm 0.031	0.243 \pm 0.031
Rainy	0.146 \pm 0.0232	0.527 \pm 0.005	0.0765 \pm 0.008
Winter	0.5682 \pm 0.0565	0.0395 \pm 0.004	0.628 \pm 0.055

The mean and stand deviation of GPP, NPP and CR of three different season are also presented in Table. Seasonally, the minimum GPP was recorded in rainy season and maximum GPP value was recorded during summer season. On monthly basis, minimum value during August and maximum value during March were recorded. The minimum value of NPP was recorded during rainy season but the maximum value was associated with summer. No particular trend was observed in seasonal variation of NPP. On comparison of the monthly variations, an increasing trend was found from the month of September to February.

Decreased value of NPP and GPP during the rainy season might be due to the fact that high suspended solids in the flood water restrict light penetration into the water and there by results in less photosynthetic activities and productivity. At the same time the phenomenon of organic matter entering the marine system, through riverine system, causing increased demand of oxygen for the oxidation of allochthonous organic matter cannot be ruled out. The higher valued of GPP and NPP during the summer season may be due to penetration of more light intensity which facilitates higher rate of photosynthesis and ultimately the productivity of the marine system.

The community respiration (CR) exhibited a systematic seasonal pattern with a maximum value during summer and maximum value during winter. A higher community respiration value during summer is possibly due to increased water temperature which stimulates the growth of microbial population which in turn utilizes more oxygen for their metabolic activities. The decreased respiration rate during winter was linked with low water temperature and reduced light which affects the rate of photosynthetic efficiency (Ahmad and Singh, 1987 and Dash *et al.*, 2011). The ratio of net and gross primary production is important for the evaluation of the amount of gross production available to the consumer (Singh and Singh, 1999). Table 1 represents the ratio between the mean and seasonal values of NPP, GPP and CR- The GPP-NPP as well as NPP-CR ratio was highest during winter and lowest in summer season while the CR percentage of GPP was found to be highest during rainy season and lowest during winter season. The NPP: CR of value > 1 during winter could be attributed to clarity of water as well as suitable temperature which favours abundance of phytoplankton and mere photosynthetic activities. During the rainy season, the ratio was < 1 , which may be due to less penetration of light into water because of suspended particles resulting in poor photosynthetic activities and thereby decrease in productivity. However, increase in community respiration, due to the flow of organic detritus along with flood water, cannot be ruled out. Higher production is not governed by a single factor as stated by Singh and Singh, (1999).

Table 2 . Mean Monthly Narration of GPP, RES & NPP values in $g^c m^3 h^{-1}$ (Mean \pm SD) at Paradip

Months	GPP	RES	NPP
January	0.670 \pm 0.055	0.036 \pm 0.005	0.788 \pm 0.057
February	0.680 \pm 0.058	0.033 \pm 0.003	0.864 \pm 0.082
March	0.930 \pm 0.105	0.180 \pm 0.018	0.787 \pm 0.068
April	0.880 \pm 0.139	0.298 \pm 0.045	0.598 \pm 0.047
May	0.740 \pm 0.066	0.198 \pm 0.025	0.572 \pm 0.055
June	0.650 \pm 0.054	0.297 \pm 0.036	0.381 \pm 0.048
July	0.290 \pm 0.030	0.078 \pm 0.008	0.139 \pm 0.007
August	0.190 \pm 0.027	0.048 \pm 0.006	0.051 \pm 0.004
September	0.076 \pm 0.008	0.037 \pm 0.003	0.031 \pm 0.005
October	0.028 \pm 0.028	0.048 \pm 0.005	0.085 \pm 0.016
November	0.396 \pm 0.065	0.058 \pm 0.006	0.372 \pm 0.036
December	0.527 \pm 0.048	0.031 \pm 0.004	0.491 \pm 0.045

There are several environmental factors acting simultaneously while must be taken into consideration while evaluating the production capacity of a marine ecosystem. There are certain physico-chemical and biological factors which in fact control the rate of production in a marine ecosystem. The paired sampled correlations among GPP, NPP and CR with other water quality parameters were carried out and the results along with significance levels were represented in table, 1, 2 and 3. The paired correlations between the variables having significant levels less than 0.05 were considered for discussion. During summer positive correlation between GPP and NPP ($r = 0.918$, $P < 0.001$) demonstrated that high GPP is responsible for high NPP. A negative correlation between GPP and EC ($r = 0.405$, $P = 0.046$) and GPP and transparency ($r = 0.463$, $p = 0.022$) was established. The above findings are in agreement with that of Ahmed *et al.*, (2005) and Dash *et al.*,

Table 3. Total & Seasonal Ratio Between Different Productivity Parameters Of Bay Of Bengal At Paradip

Total			Seasonal								
NPP:GPP	NPP:CR	CR% Of GPP	NPP: GPP			NPP : CR			CR % of GPP		
			S	R	W	S	R	W	S	R	W
0.84	3.95	20.44	0.74	0.46	1.15	2.49	1.89	16.21	28.8	26.67	6.51

(2011) where they reported a moderate correlation between EC and GPP and NPP, A positive relationship between GPP and BOD ($r = 0.425$, $p = 0.005$) indirectly states that the growth of phytoplankton and their increased productivity is due to organic wastes, containing high nutrient values. In summer season, BOD plays a major role in controlling (GPP, NPP and CR. Positive relationships between GPP with CR ($r = 0.655$, $P < 0.001$) and GPP with NPP ($r = 0.761$, $P < 0.001$) were observed. Higher alkalinity and hardness are favourable for increase in GPP and NPP during rainy season. The findings of Goldman and Wetzel (1963), Singh (1999) and Dash *et al.* (2011), support the above finding, positive correlation between alkalinity and GPP, during rainy season was also reported by Yeragi and Shaikh, (2003) and Dash *et al.*, 2011)

During winter a negative relationship between GPP and CR ($r = -0.53$, $p = 0.005$) and GPP with water temperature ($r = -0.497$, $P = 0.013$) was noted in the current study. An inverse relationship between GPP and alkalinity and NPP with water temperature was recorded. Nitrate being the most essential nutrient for the growth of aquatic organism supports the positive relation between CR and nitrate during this season.

Acknowledgement

The author is grateful to Head of the P.G Dept. of Zoology, Vani Vihar, Utkal University, BBSR, Odisha for providing necessary laboratory facilities to carry out this work and valuable suggestion to carry out the work.

REFERENCES

Ahmed, S.H. and Singh, A.K., 1987. Seasonal, fluctuations of primary production and fish yields in mangrove's tank, Patna City (Bihar). *Geobios*, 14:62-66.

- Ahmed, K.K.U., Ahmed, S.V., Halder, G.C., Hossain, M.R.A., and Ahmed, T.S., 2005. Primary production and fish yield estimation in the meghna river system, Bangladesh. *Asian Fisheries Science*, 18:95-105.
- APHA, 1998. American water works Association and water pollution control Federation standard methods for examination of water and waste water. *Am. Pub. Health. Assoc.*, New York, USA.
- Dash, S., Patra, A.K. and Adhikari, S., 2011. Primary Productivity of Kharasrota river (India) *J. Ecophysiol. Occup. Hlth.*, 11:219-225.
- Gaarder T. and Gran H.H., 1927. Investigation on the production of plankton in the Oslo Fjord, P-v Reun. *Commn. Inter explor Sci, Mer. Medieterr.*, 42:1-48.
- Goldman, C.R. and Wetzel, R.G., 1963. A study of the Primary productivity of clear lake, Lake Country, California, *Ecology*, 44:283-294.
- Odum E.P., 1971. Fundamentals of Ecology, 3rd edn. W.B. Saunders Company, Philadelphia:574.
- Odum, H.T., 1956. Primary Production in flowing waters. *Limo. Oceanogr*, 1: 102-117.
- Singh, A.K. and Singh, D.K., 1999. A comparative study of the phytoplanktonic primary production of river Ganga and Pond of Patna (Bihar), *India. J. Environ. Biol.*, 20:263-270.
- Thomas, P. A., Abraham, T. and Abraham, K.G., 1980. Observation on the Primary productivity of Sasthamkotta lake: Dn. Alexander, K.M. et al (eds) *Proe. Symp. Environ. Biol. Trivandram*: 1-7.
- Vollenweider, R.A., 1969. A manual on methods for measuring production in the aquatic environment IBP, Hand Book No.12, Black well scientific publications, Oxford, England, 138:465-482.
- Yeragi, S.G. and Shaikh, N. 2003. Studies on Primary productivity of Tansa river. *J. Natcon.*, 15: 125 - 130.
